



National Comprehensive
Cancer Network®

NCCN Clinical Practice Guidelines in Oncology (NCCN Guidelines®)

Chronic Myeloid Leukemia

Version 2.2022 — November 15, 2021

NCCN.org

NCCN Guidelines for Patients® available at www.nccn.org/patients

Continue



National
Comprehensive
Cancer
Network®

NCCN Guidelines Version 2.2022

Chronic Myeloid Leukemia

[NCCN Guidelines Index](#)
[Table of Contents](#)
[Discussion](#)

Neil P. Shah, MD, PhD/Chair ‡
UCSF Helen Diller Family
Comprehensive Cancer Center

Ravi Bhatia, MD/Vice-Chair ‡
O'Neal Comprehensive
Cancer Center at UAB

Jessica K. Altman, MD ‡
Robert H. Lurie Comprehensive Cancer
Center of Northwestern University

Maria Amaya, MD, PhD ‡
University of Colorado Cancer Center

Ellin Berman, MD ‡ † P
Memorial Sloan Kettering Cancer Center

Robert H. Collins, Jr., MD ‡
UT Southwestern Simmons
Comprehensive Cancer Center

Peter T. Curtin, MD † ‡ ξ
City of Hope National Medical Center

Daniel J. DeAngelo, MD, PhD ‡ †
Dana-Farber/Brigham and Women's
Cancer Center

Jason Gotlib, MD, MS ‡ †
Stanford Cancer Institute

Gabriela Hobbs, MD ‡ †
Massachusetts General Hospital
Cancer Center

Lori Maness, MD ‡
Fred and Pamela Buffett Cancer Center

Monica Mead, MD †
UCLA Jonsson Comprehensive Cancer Center

Leland Metheny, MD ‡ ξ
Case Comprehensive Cancer Center
University Hospitals Seidman Cancer Center and
Cleveland Clinic Taussig Cancer Institute

Sanjay Mohan, MD, MSCI ‡
Vanderbilt-Ingram Cancer Center

Joseph O. Moore, MD †
Duke Cancer Institute

Vivian Oehler, MD ‡
Fred Hutchinson Cancer Research Center/
Seattle Cancer Care Alliance

Mrinal Patnaik, MD †
Mayo Clinic Cancer Center

Keith Pratz, MD †
Abramson Cancer Center
at the University of Pennsylvania

Iskra Pusic, MD, MSCI †
Siteman Cancer Center at Barnes-
Jewish Hospital and Washington
University School of Medicine

Michal G. Rose, MD †
Yale Cancer Center/
Smilow Cancer Hospital

B. Douglas Smith, MD † P
The Sidney Kimmel Comprehensive
Cancer Center at Johns Hopkins

Kendra L. Sweet, MD, MS ‡ † P
Moffitt Cancer Center

Moshe Talpaz, MD †
University of Michigan
Rogel Cancer Center

Tiffany N. Tanaka, MD ‡
UC San Diego Moores Cancer Center

James Thompson, MD, MS ‡
Roswell Park Comprehensive
Cancer Center

Jennifer Vaughn, MD, MSPH ‡
The Ohio State University Comprehensive
Cancer Center - James Cancer Hospital
and Solove Research Institute

Jeanna Welborn, MD †
UC Davis Comprehensive Cancer Center

David T. Yang, MD ≠
University of Wisconsin
Carbone Cancer Center

NCCN
Kristina Gregory, RN, MSN, OCN
Hema Sundar, PhD

ξ Bone marrow transplantation
‡ Hematology/Hematology oncology
P Internal medicine
† Medical oncology
≠ Pathology

[NCCN Guidelines Panel Disclosures](#)

Continue



National
Comprehensive
Cancer
Network®

NCCN Guidelines Version 2.2022

Chronic Myeloid Leukemia

[NCCN Guidelines Index](#)
[Table of Contents](#)
[Discussion](#)

[NCCN Chronic Myeloid Leukemia Panel Members](#)
[Summary of Guidelines Updates](#)

[Workup \(CML-1\)](#)

[Chronic Phase CML \(CML-2\)](#)

[Early Treatment Response Milestones, Clinical Considerations, and Recommendations \(CML-3\)](#)

[Advanced Phase CML \(CML-4\)](#)

[Treatment Recommendations Based on *BCR-ABL1* Mutation Profile \(CML-5\)](#)

[Allogeneic Hematopoietic Cell Transplantation \(CML-6\)](#)

[Risk Calculation Table \(CML-A\)](#)

[Definitions of Accelerated Phase and Blast Phase \(CML-B\)](#)

[Management of CML During Pregnancy \(CML-C\)](#)

[Criteria for Response and Relapse \(CML-D\)](#)

[Monitoring Response to TKI Therapy and Mutational Analysis \(CML-E\)](#)

[Discontinuation of TKI Therapy \(CML-F\)](#)

[Management of Bosutinib Toxicity \(CML-G 1 of 9\)](#)

[Management of Dasatinib Toxicity \(CML-G 2 of 9\)](#)

[Management of Imatinib Toxicity \(CML-G 3 of 9\)](#)

[Management of Nilotinib Toxicity \(CML-G 4 of 9\)](#)

[Management of Omacetaxine Toxicity \(CML-G 5 of 9\)](#)

[Management of Ponatinib Toxicity \(CML-G 6 of 9\)](#)

[Management of Asciminib Toxicity \(CML-G 7 of 9\)](#)

[Drug Interactions of TKIs \(CML-G 8 of 9\)](#)

Clinical Trials: NCCN believes that the best management for any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.

Find an NCCN Member Institution:
<https://www.nccn.org/home/member-institutions>.

NCCN Categories of Evidence and Consensus: All recommendations are category 2A unless otherwise indicated.

See [NCCN Categories of Evidence and Consensus](#).

NCCN Categories of Preference: All recommendations are considered appropriate.

See [NCCN Categories of Preference](#).

The NCCN Guidelines® are a statement of evidence and consensus of the authors regarding their views of currently accepted approaches to treatment. Any clinician seeking to apply or consult the NCCN Guidelines is expected to use independent medical judgment in the context of individual clinical circumstances to determine any patient's care or treatment. The National Comprehensive Cancer Network® (NCCN®) makes no representations or warranties of any kind regarding their content, use or application and disclaims any responsibility for their application or use in any way. The NCCN Guidelines are copyrighted by National Comprehensive Cancer Network®. All rights reserved. The NCCN Guidelines and the illustrations herein may not be reproduced in any form without the express written permission of NCCN. ©2021.



NCCN Guidelines Version 2.2022

Chronic Myeloid Leukemia

Updates in Version 2.2022 of the NCCN Guidelines for Chronic Myeloid Leukemia from Version 1.2022 include:

[CML-5](#)

- Asciminib added to the Table for Treatment Recommendations Based on *BCR-ABL1* Mutation Profile
- Footnote x added: Asciminib is a treatment option for CP-CML patients with the T315I mutation and/or CP-CML with resistance or intolerance to at least two prior TKIs.

[CML-F](#)

- Footnote 1 modified: The feasibility of TFR following discontinuation of ~~bosutinib or ponatinib~~ *TKIs other than dasatinib, imatinib, or nilotinib* has not yet been evaluated in clinical studies.

[CML-G 6 of 9](#)

- Bullet one modified: ~~Vascular occlusion: Arterial and venous thrombosis and occlusions, including fatal myocardial infarction and stroke, have occurred at a considerable rate~~ *Arterial occlusive events (AOEs; including fatal myocardial infarction and stroke) and venous thromboembolic events (VTEs), have occurred* in patients treated with ponatinib.

[CML-G 7 of 9](#)

- New page added for the Management of Asciminib Toxicity



NCCN Guidelines Version 2.2022

Chronic Myeloid Leukemia

Updates in Version 1.2022 of the NCCN Guidelines for Chronic Myeloid Leukemia from Version 3.2021 include:

General: Terminologies modified to be more inclusive of sexual and gender identities.

[CML-2](#)

- Primary Treatment; Imatinib
 - ▶ Generic wording removed (also applies to CML-4)
- Footnote f modified: *If treatment is needed during pregnancy, it is preferable to initiate treatment with interferons (interferon alfa-2a or peginterferon alfa-2a). Interferon alfa-2a/2b and peginterferon alfa-2b have been discontinued. Peginterferon alfa-2a may be substituted for other interferon preparations. TKI therapy, particularly during the first trimester, should be avoided. See Management of CML During Pregnancy (CML-C).*
- Footnote g modified: Based on follow-up data from the BFORE, DASISION, and ENESTnd trials, second-generation TKIs (bosutinib, dasatinib, or nilotinib) are preferred for patients with an intermediate- or high-risk score. *Second-generation TKIs should also be considered for specific subgroups (based on the assessment of treatment goals and benefit/risks), for example, younger patients who are interested in ultimately discontinuing treatment and especially for young women patients assigned female at birth whose goal is to achieve a deep and rapid molecular response and eventual drug discontinuation of TKI therapy for family planning purposes.*
- Footnote h added: Innovator and generic drugs approved by the regulatory authorities based on pharmacokinetic equivalence can be used interchangeably. An FDA-approved generic version is an appropriate substitute for an innovator drug (imatinib). Generic versions of other TKIs are likely to be marketed in the near future. (also applies to CML-4, CML-6)

[CML-3](#)

- Clinical Considerations; Light Green, TKI-sensitive disease
 - ▶ If treatment goal is long-term survival: $\geq 0.4\%$ – $\leq 1\%$ optimal

[CML-5](#)

- Nilotinib: G250E removed as a contraindicated mutation

[CML-6](#)

- Not in CCyR or in relapse changed to Less than CCyR or in relapse
- Follow-up Therapy; PCR, Positive
 - ▶ Discuss options with transplant team: TKI $\pm$ DLI (selection of TKI is based on prior therapy depending on prior TKI, tolerance, BCR-ABL1 mutation profile, and post-HCT morbidities)

- Footnote aa changed from a link to the Discussion to the following references: Carpenter PA, et al. Blood 2007;109:2791-2793; Olavarria E, et al. Blood 2007;110:4614-4617; DeFilipp Z, et al. Clin Lymphoma Myeloma Leuk 2016;16:466-471.

[CML-C 1 of 2](#)

- Treatment and Monitoring During Pregnancy, Bullet 4 modified:
 - ▶ *If treatment is needed, it is preferable to initiate treatment with interferons. Both interferon alfa-2a or peginterferon alfa-2a have been used during pregnancy. Most of the data using interferons during pregnancy have been reported in patients with essential thrombocythemia. If introduced earlier, the use of interferon alfa-2a/2b or peginterferon alfa-2a can preserve molecular remission after discontinuation of TKI. Interferon alfa-2a/2b and peginterferon alfa-2b have been discontinued. Peginterferon alfa-2a may be substituted for other interferon preparations. Data are insufficient to establish the use of peginterferon alfa-2a (risk category C) in pregnancy and it should be used only if benefits outweigh potential risk to the fetus.*

[CML-C 2 of 2](#)

- Reference 15 added.


[CML-D](#)

- Title modified: Criteria for Hematologic, Cytogenetic, and Molecular Response and Relapse
- Relapse definition modified (also applies to CML-E)
 - ▶ ~~Any sign of loss of response (defined as hematologic or cytogenetic relapse)~~
 - ▶ *Any sign of loss of hematologic response*
 - ▶ *Any sign of loss of CCyR or its molecular response correlate defined as an increase in BCR-ABL1 transcript to $>1\%$*
 - ▶ *1-log increase in BCR-ABL1 transcript levels with loss of MMR should prompt bone marrow evaluation for loss of CCyR but is not itself defined as relapse (eg, hematologic or cytogenetic relapse)*
- Footnote 8 added: The loss of MMR in the presence of a CCyR does not necessarily constitute treatment failure.



NCCN Guidelines Version 2.2022

Chronic Myeloid Leukemia



^a Bone marrow evaluation should be done for the initial workup, to provide morphologic review, and also to detect chromosomal abnormalities in addition to the Ph chromosome. Fluorescence in situ hybridization (FISH) can be used if cytogenetic evaluation is not possible.

^b Consider qualitative RT-PCR for the detection of atypical *BCR-ABL1* transcripts. See [Discussion](#). Referral to centers with expertise in the management of rare hematologic malignancies is recommended.

^c Hepatitis B virus reactivation has been reported in patients receiving TKI therapy. However, it is not always possible to reliably estimate the frequency or establish a relationship to drug exposure because these incidences are reported voluntarily from a population of uncertain size.

^d See [Definitions of Accelerated Phase and Blast Phase \(CML-B\)](#).

^e Consider myeloid mutation panel for patients with accelerated phase or blast phase.

Note: All recommendations are category 2A unless otherwise indicated.


Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.



NCCN Guidelines Version 2.2022

Chronic Myeloid Leukemia

CLINICAL PRESENTATION



^f If treatment is needed during pregnancy, it is preferable to initiate treatment with interferons (interferon alfa-2a or peginterferon alfa-2a). Interferon alfa 2a/2b and peginterferon alfa-2b have been discontinued. Peginterferon alfa-2a may be substituted for other interferon preparations. TKI therapy, particularly during the first trimester, should be avoided. [See Management of CML During Pregnancy \(CML-C\)](#).

⁹ Based on follow-up data from the BFORE, DASISION, and ENESTnd trials, second-generation TKIs (bosutinib, dasatinib, or nilotinib) are preferred for patients with an intermediate- or high-risk score. Second-generation TKIs should also be considered for specific subgroups (based on the assessment of treatment goals and benefit/risks), for example, younger patients who are interested in ultimately discontinuing treatment and especially young patients assigned female at birth whose goal is to achieve a deep and rapid molecular response and eventual discontinuation of TKI therapy for family planning purposes.

^h Innovator and generic drugs approved by the regulatory authorities based on pharmacokinetic equivalence can be used interchangeably. An FDA-approved generic version is an appropriate substitute for an innovator drug (imatinib). Generic versions of other TKIs are likely to be marketed in the near future.

ⁱ Imatinib may be preferred for older patients with comorbidities such as cardiovascular disease.

^j [See Criteria for Response and Relapse \(CML-D\)](#).

^k [See Monitoring Response to TKI Therapy and Mutational Analysis \(CML-E\)](#).

Note: All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.



NCCN Guidelines Version 2.2022

Chronic Myeloid Leukemia

EARLY TREATMENT RESPONSE MILESTONES^{j,k}

<i>BCR-ABL1</i> (IS)	3 months	6 months	12 months ^l
>10% ^m	YELLOW	RED	
>1%–10%	GREEN		YELLOW
>0.1%–1%	GREEN		LIGHT GREEN
≤0.1%	GREEN		

COLOR	CONCERN	CLINICAL CONSIDERATIONS	RECOMMENDATIONS
RED	TKI-resistant disease	- Evaluate patient compliance and drug interactions - Consider mutational analysis	Switch to alternate TKI (CML-5) and evaluate for allogeneic HCT
YELLOW	Possible TKI resistance	- Evaluate patient compliance and drug interactions - Consider mutational analysis - Consider bone marrow cytogenetic analysis to assess for MCyR at 3 mo or CCyR at 12 mo	Switch to alternate TKI (CML-5) or Continue same TKI (other than imatinib) (CML-G) ⁿ or Increase imatinib dose to a max of 800 mg and Consider evaluation for allogeneic HCT
LIGHT GREEN	TKI-sensitive disease	- If treatment goal is long-term survival: ≤1% optimal - If treatment goal is treatment-free remission: ≤0.1% optimal	- If optimal: continue same TKI - If not optimal: shared decision-making with patient^{o,p}
GREEN	TKI-sensitive disease	Monitor response (CML-E) and side effects	Continue same TKI (CML-G) ^q

^j See [Criteria for Response and Relapse \(CML-D\)](#).

^k See [Monitoring Response to TKI Therapy and Mutational Analysis \(CML-E\)](#).

^l *BCR-ABL1* ≤0.1% at 12 months is associated with a very low probability of subsequent loss of response and a high likelihood of achieving a subsequent deep molecular response (MR4.0; ≤0.01% *BCR-ABL1* IS), which is a prerequisite for a trial of treatment-free remission (TFR).

^m Patients with *BCR-ABL1* only slightly >10% at 3 months and/or with a steep decline from baseline may achieve <10% at 6 months and have generally favorable outcomes. Therefore, it is important to interpret the value at 3 months in this context before making drastic changes to the treatment strategy.

ⁿ Achievement of response milestones must be interpreted within the clinical context. Patients with more than 50% reduction compared to baseline or minimally above the 10% cutoff can continue the same dose of dasatinib, nilotinib, or bosutinib for another 3 months. Continuation of imatinib 400 mg is not recommended.

^o Switching from imatinib to a second-generation TKI improves response, but is associated with increased toxicity.

^p Consider referral to a specialized CML center and/or enrollment in a clinical trial.

^q Discontinuation of TKI with careful monitoring is feasible in selected patients. See [Discontinuation of TKI Therapy \(CML-F\)](#).

Note: All recommendations are category 2A unless otherwise indicated.


Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.



NCCN Guidelines Version 2.2022

Chronic Myeloid Leukemia

CLINICAL PRESENTATION



^d [See Definitions of Accelerated Phase and Blast Phase \(CML-B\)](#).

^f If treatment is needed during pregnancy, it is preferable to initiate treatment with interferons (interferon alfa-2a or peginterferon alfa-2a). Interferon alfa 2a/2b and peginterferon alfa-2b have been discontinued. Peginterferon alfa-2a may be substituted for other interferon preparations. TKI therapy, particularly during the first trimester, should be avoided. [See Management of CML During Pregnancy \(CML-C\)](#).

^h Innovator and generic drugs approved by the regulatory authorities based on pharmacokinetic equivalence can be used interchangeably. An FDA-approved generic version is an appropriate substitute for innovator drug (imatinib). Generic versions of other TKIs are likely to be marketed in the near future.

^r Consider evaluation for allogeneic HCT if response milestones are not achieved at 3, 6, and 12 months as outlined on [CML-3](#).

^s The presence of major route additional chromosomal abnormalities in Ph-positive cells (ACA/Ph+; trisomy 8, isochromosome 17q, second Ph, and trisomy 19) may have a negative prognostic impact on survival. Patients who present with accelerated phase at diagnosis should be treated with a TKI, followed by evaluation for allogeneic HCT, based on response to therapy.

^t Imatinib is not recommended for patients with disease progression on prior TKI therapy.

^u Omacetaxine is a treatment option for patients with disease progression to accelerated phase CML. Omacetaxine is not a treatment option for patients who present with accelerated phase CML.

Note: All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.



NCCN Guidelines Version 2.2022

Chronic Myeloid Leukemia

TREATMENT RECOMMENDATIONS BASED ON *BCR-ABL1* MUTATION PROFILE

- Patients with disease resistant to primary treatment with imatinib should be treated with bosutinib, dasatinib, or nilotinib in the second-line setting, taking into account *BCR-ABL1* mutation status.
- Patients with disease resistant to primary treatment with bosutinib, dasatinib, or nilotinib can be treated with an alternate TKI (other than imatinib) in the second-line setting, taking into account *BCR-ABL1* mutation status. The durability of these responses is frequently limited.
- The table below lists the *BCR-ABL1* mutations that should NOT be treated with bosutinib, dasatinib, or nilotinib in the second-line setting.

THERAPY	CONTRAINDICATED MUTATIONS ^v
Bosutinib	T315I, V299L, G250E, or F317L ^w
Dasatinib	T315I/A, F317L/V/I/C, or V299L
Nilotinib	T315I, Y253H, E255K/V, or F359V/C/I
Asciminib , ^x Ponatinib , ^y Omacetaxine , ^z allogeneic HCT (CML-6), or clinical trial	None

^v Mutations contraindicated for imatinib are too numerous to include. There are compound mutations that can cause resistance to ponatinib, but those are uncommon following treatment with bosutinib, dasatinib, or nilotinib.

^w Bosutinib has minimal activity against F317L mutation. Nilotinib may be preferred over bosutinib in patients with an F317L mutation.

^x Asciminib is a treatment option for CP-CML patients with the T315I mutation and/or CP-CML with resistance or intolerance to at least two prior TKIs.

^y Ponatinib is a treatment option for patients with a T315I mutation and/or CP-CML with resistance or intolerance to at least two prior TKIs or for patients with AP-CML or BP-CML for whom no other TKI is indicated.


^z Omacetaxine is a treatment option for patients with disease that is resistant and/or intolerant to two or more TKIs.

Note: All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.



FOLLOW-UP THERAPY



^h Innovator and generic drugs approved by the regulatory authorities based on pharmacokinetic equivalence can be used interchangeably. An FDA-approved generic version is an appropriate substitute for innovator drug (imatinib). Generic versions of other TKIs are likely to be marketed in the near future.

^j [See Criteria for Response and Relapse \(CML-D\).](#)

^{aa} Indications for allogeneic HCT: advanced phase CML at presentation or disease progression to blast phase. Outcomes of allogeneic HCT are dependent on age and comorbidities, donor type, and transplant center.

^{bb} Carpenter PA, et al. Blood 2007;109:2791-2793; Olavarria E, et al. Blood 2007;110:4614-4617; DeFilipp Z, et al. Clin Lymphoma Myeloma Leuk 2016;16:466-471.

Note: All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.



NCCN Guidelines Version 2.2022

Chronic Myeloid Leukemia

RISK CALCULATION TABLE

Risk Score	Calculation	Risk Category
Sokal score¹	$\text{Exp } 0.0116 \times (\text{age} - 43.4) + 0.0345 \times (\text{spleen} - 7.51) + 0.188 \times [(\text{platelet count} \div 700)^2 - 0.563] + 0.0887 \times (\text{blasts} - 2.10)$	Low <0.8 Intermediate 0.8 – 1.2 High >1.2
Hasford (EURO) score²	$(0.6666 \times \text{age} [0 \text{ when age} < 50 \text{ years}; 1, \text{ otherwise}] + 0.042 \times \text{spleen size [cm below costal margin]} + 0.0584 \times \text{percent blasts} + 0.0413 \times \text{percent eosinophils} + 0.2039 \times \text{basophils} [0 \text{ when basophils} < 3\%; 1, \text{ otherwise}] + 1.0956 \times \text{platelet count} [0 \text{ when platelets} < 1500 \times 10^9/\text{L}; 1, \text{ otherwise}]) \times 1000$	Low ≤780 Intermediate >780 – ≤1480 High >1480
EUTOS long-term survival (ELTS) score³	$0.0025 \times (\text{age}/10)^3 + 0.0615 \times \text{spleen size cm below costal margin} + 0.1052 \times \text{blasts in peripheral blood} + 0.4104 \times (\text{platelet count}/1000)^{-0.5}$	Low ≤1.5680 Intermediate >1.5680 but ≤2.2185 High >2.2185

Calculation of relative risk based on Sokal or Hasford (EURO) score can be found at:

https://www.leukemia-net.org/content/leukemias/cml/euro_and_sokal_score/index_eng.html

Online calculator for the ELTS score can be found at: https://www.leukemia-net.org/content/leukemias/cml/elts_score/index_eng.html

¹ Sokal J, Cox EB, Baccarani M, et al. Prognostic discrimination in "good-risk" chronic granulocytic leukemia. Blood 1984;63:789-799.

² Hasford J, Pfirrmann M, Hehlmann R, et al. A new prognostic score for survival of patients with chronic myeloid leukemia treated with interferon alfa. Writing Committee for the Collaborative CML Prognostic Factors Project Group. J Natl Cancer Inst 1998;90:850-858.

³ Pfirrmann M, Baccarani M, Saussele S, et al. Prognosis of long-term survival considering disease-specific death in patients with chronic myeloid leukemia. Leukemia 2016;30:48-56.

Note: All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.



DEFINITIONS OF ACCELERATED PHASE^{1,2}

Modified MD Anderson Cancer Center (MDACC) Criteria^{3,4} (most commonly used in clinical trials)

- Peripheral blood myeloblasts $\geq 15\%$ and $< 30\%$
- Peripheral blood myeloblasts and promyelocytes combined $\geq 30\%$
- Peripheral blood basophils $\geq 20\%$
- Platelet count $\leq 100 \times 10^9/L$ unrelated to therapy
- Additional clonal cytogenetic abnormalities in Ph+ cells⁵

DEFINITIONS OF BLAST PHASE¹

International Bone Marrow Transplant Registry^{6,7}

- $\geq 30\%$ blasts in the blood, marrow, or both
- Extramedullary infiltrates of leukemic cells

¹ Any increase in lymphoblasts is concerning for (nascent) blast phase.

² Sokal criteria (Sokal JE, Baccarani M, Russo D, Tura S. Staging and prognosis in chronic myelogenous leukemia. Semin Hematol 1988;25:49-61) and IBMTR criteria (Savage DG, Szydlo RM, Chase A, et al. Bone marrow transplantation for chronic myeloid leukemia: The effects of differing criteria for defining chronic phase on probabilities of survival and relapse. Br J Haematol 1997;99:30-35) are historically used when HCT is the recommended treatment option.

³ Kantarjian HM, Deisseroth A, Kurzrock R, et al. Chronic myelogenous leukemia: A concise update. Blood 1993;82:691-703.

⁴ Talpaz M, Silver RT, Druker BJ, et al. Imatinib induces durable hematologic and cytogenetic responses in patients with accelerated phase chronic myeloid leukemia: results of a phase 2 study. Blood 2002;99:1928-1937.

⁵ The prognostic significance of additional chromosomal abnormalities in Ph-positive cells (ACA/Ph+) is related to the specific chromosomal abnormality and often other features of accelerated phase. The presence of "major route" ACA/Ph+ (trisomy 8, isochromosome 17q, second Ph, and trisomy 19) at diagnosis may have a negative prognostic impact on survival.

⁶ Druker BJ. Chronic Myelogenous Leukemia In: DeVita VT, Lawrence TS, Rosenberg SA, eds. DeVita, Hellman, and Rosenberg's Cancer: Principles & Practice of Oncology. Vol. 2 (ed 8): Lippincott, Williams and Wilkins; 2007:2267-2304.

⁷ World Health Organization (WHO) criteria may be included in some reports (Swerdlow SH, Harris NL, Jaffe ES, et al. WHO Classification of Tumours of Haematopoietic and Lymphoid Tissues. Revised 4th ed. Lyon, France: IARC; 2017): blasts $\geq 20\%$ of peripheral white blood cells or of nucleated bone marrow cells; extramedullary blast proliferation; and large foci or clusters of blasts in the bone marrow biopsy. However, it should be noted that modified MDACC criteria were used in most clinical trials leading to the approval of TKIs.

Note: All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.

**MANAGEMENT OF CML DURING PREGNANCY****Tyrosine Kinase Inhibitor (TKI) Therapy and Conception**

- TKI therapy appears to affect some male hormones at least transiently, but does not appear to have a deleterious effect on male fertility; miscarriage or fetal abnormality rate is not elevated in female partners of male patients on TKI therapy.¹⁻⁵
- TKI therapy during pregnancy in patients assigned female at birth has been associated with both a higher rate of miscarriage and fetal abnormalities. A prolonged washout period prior to pregnancy, prompt consideration of holding TKI therapy (if pregnancy occurs while on TKI therapy), and close monitoring should be considered.⁶⁻¹⁰
- Discontinuation of TKI therapy because of pregnancy in patients who were not in deep molecular response (DMR $\geq$ MR4.0; $\leq$ 0.01% *BCR ABL1* IS) has only been reported in a small series of patients.¹¹⁻¹⁴ Conception while on active TKI therapy is strongly discouraged due to the risk of fetal abnormalities.
- Prior to attempting pregnancy, patients of childbearing age and their partners should be counseled about the potential risks and benefits of discontinuation of TKI therapy and possible resumption of TKI therapy should CML recur during pregnancy. Fertility preservation should be discussed with all patients of childbearing age prior to the initiation of TKI therapy. Referral to a CML specialty center and consultation with a high-risk obstetrician is recommended.

Treatment and Monitoring During Pregnancy

- In patients assigned male at birth, TKI therapy need not be discontinued if a pregnancy is planned. Sperm banking can also be performed prior to starting TKI therapy, although there are no data regarding the quality of sperm in patients with untreated CML.
- In patients assigned female at birth, TKI therapy should be stopped prior to natural conception, and patients should remain off therapy during pregnancy.⁶⁻⁸ Referral to an in vitro fertilization (IVF) center is recommended in coordination with the patient's obstetrician. TKI should be stopped prior to attempting a natural pregnancy or oocyte retrieval, but the optimal timing of discontinuation is unknown.
- The use of TKI therapy, particularly during the first trimester, should be avoided. If TKI therapy is considered during pregnancy, the potential risks and benefits must be carefully evaluated in terms of maternal health and fetal risk on an individual basis prior to initiation of TKI therapy during pregnancy.

- If treatment is needed, it is preferable to initiate treatment with interferons. Both interferon alfa-2a or peginterferon alfa-2a have been used during pregnancy.¹⁵⁻²¹ Most of the data using interferons during pregnancy have been reported in patients with essential thrombocythemia.^{15,21} If introduced earlier, the use of interferon alfa-2a or peginterferon alfa-2a can preserve molecular remission after discontinuation of TKI.²² Interferon alfa-2a/2b and peginterferon alfa-2b have been discontinued. Peginterferon alfa-2a may be substituted for other interferon preparations.
- The panel recommends against the use of hydroxyurea during pregnancy, especially in the first trimester, if possible.²³⁻²⁵
- Leukapheresis can be used for a rising white blood cell (WBC) count, although there are no data that recommend at what level of WBC count leukapheresis should be initiated.²⁶⁻²⁹
- Low-dose aspirin or low-molecular-weight heparin can be considered for patients with thrombocytosis.^{30,31}
- Monthly monitoring with qPCR and initiating treatment if the *BCR ABL1* IS increases to $>1.0\%$ is recommended.

Breastfeeding

- TKI therapy can be restarted after delivery. However, patients should be advised not to breastfeed while on TKI therapy, as TKIs pass into human breast milk.³²⁻³⁵
- Breastfeeding without TKI therapy may be safe with molecular monitoring, but preferably in those patients with CML who have achieved durable DMR. It may be acceptable to avoid TKIs for the short period of the first 2–5 days after labor to give the child colostrum.^{35,36}
- Close molecular monitoring is recommended for patients who extend the treatment-free period for breastfeeding. If the loss of MMR after treatment cessation is confirmed, breastfeeding needs to be terminated and TKI therapy should be restarted.³⁵

Note: All recommendations are category 2A unless otherwise indicated.**Clinical Trials:** NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.[References](#)
[\(CML-C 2 of 2\)](#)**CML-C**
1 OF 2



MANAGEMENT OF CML DURING PREGNANCY – REFERENCES

- ¹ Ramasamy K, Hayden J, Lim Z, et al. Successful pregnancies involving men with chronic myeloid leukaemia on imatinib therapy. *Br J Haematol* 2007;137:374-375.
- ² Breccia M, Cannella L, Montefusco E, et al. Male patients with chronic myeloid leukemia treated with imatinib involved in healthy pregnancies: report of five cases. *Leuk Res* 2008;32:519-520.
- ³ Oweini H, Otrrock ZK, Mahfouz RAR, Bazarbachi A. Successful pregnancy involving a man with chronic myeloid leukemia on dasatinib. *Arch Gynecol Obstet* 2011;283:133-134.
- ⁴ Ghalaut VS, Prakash G, Bansal P, et al. Effect of imatinib on male reproductive hormones in BCR-ABL positive CML patients: A preliminary report. *J Oncol Pharm Pract* 2014;20:243-248.
- ⁵ Alizadeh H, Jaafar H, Rajnics P, et al. Outcome of pregnancy in chronic myeloid leukaemia patients treated with tyrosine kinase inhibitors: short report from a single centre. *Leuk Res* 2015;39:47-51.
- ⁶ Pye SM, Cortes J, Ault P, et al. The effects of imatinib on pregnancy outcome. *Blood* 2008;111:5505-5508.
- ⁷ Cortes JE, Abruzzese E, Chelysheva E, et al. The impact of dasatinib on pregnancy outcomes. *Am J Hematol* 2015;90:1111-1115.
- ⁸ Barkoulas T, Hall PD. Experience with dasatinib and nilotinib use in pregnancy. *J Oncol Pharm Pract* 2018;24:121-128.
- ⁹ Salem W, Li K, Krapp C, et al. Imatinib treatments have long-term impact on placentation and embryo survival. *Sci Rep* 2019;9:2535.
- ¹⁰ Madabhavi I, Sarkar M, Modi M, Kadakol N. Pregnancy outcomes in chronic myeloid leukemia: A single center experience. *J Glob Oncol* 2019;5:1-11.
- ¹¹ Ault P, Kantarjian H, O'Brien S, et al. Pregnancy among patients with chronic myeloid leukemia treated with imatinib. *J Clin Oncol* 2006;24:1204-1208.
- ¹² Kuwabara A, Babb A, Ibrahim A, et al. Poor outcome after reintroduction of imatinib in patients with chronic myeloid leukemia who interrupt therapy on account of pregnancy without having achieved an optimal response. *Blood* 2010;116:1014-1016.
- ¹³ Lasica M, Willcox A, Burbury K, et al. The effect of tyrosine kinase inhibitor interruption and interferon use on pregnancy outcomes and long-term disease control in chronic myeloid leukemia. *Leuk Lymphoma* 2019;60:1796-1802:4-7.
- ¹⁴ Stella S, Tirro E, Massimino M, et al. Successful management of a pregnant patient with chronic myeloid leukemia receiving standard dose imatinib. *In Vivo* 2019;33:1593-1598.
- ¹⁵ Schrickel L, Heideel FH, Sadjadian P et al. Interferon alpha for essential thrombocythemia during 34 high-risk pregnancies: outcome and safety. *J Cancer Res and Clin Oncol* 2021; 147;1481-1491.
- ¹⁶ Haggstrom J, Adriansson M, Hybbinette T, et al. Two cases of CML treated with alpha-interferon during second and third trimester of pregnancy with analysis of the drug in the newborn immediately postpartum. *Eur J Haematol* 1996;57:101-102.
- ¹⁷ Kuroiwa M, Gondo H, Ashida K, et al. Interferon-alpha therapy for chronic myelogenous leukemia during pregnancy. *Am J Hematol* 1998;59:101-102.
- ¹⁸ Lipton JH, Derzko CM, Curtis J. Alpha-interferon and pregnancy in a patient with CML. *Hematol Oncol* 1996;14:119-122.
- ¹⁹ Al Bahar S, Pandita R, Nath SV. Pregnancy in chronic myeloid leukemia patients treated with alpha interferon. *Int J Gynaecol Obstet* 2004;85:281-282.
- ²⁰ Balsat M, Etienne M, Elhamri M, et al. Successful pregnancies in patients with BCR-ABL-positive leukemias treated with interferon-alpha therapy during the tyrosine kinase inhibitors era. *Eur J Haematol* 2018;101:774-780.
- ²¹ Beauverd Y, Radia D, Cargo C, et al. Pegylated interferon alpha-2a for essential thrombocythemia during pregnancy: outcome and safety. A case series. *Haematologica* 2016;101:e182-184.
- ²² Abruzzese E, Turkina AG, Apperley JF, et al. Pregnancy management in CML patients: To treat or not to treat? Report of 224 outcomes of the European Leukemia Net (ELN) Database [abstract]. *Blood* 2019;134:Abstract 498.
- ²³ Baykal C, Zengin N, Coskun F, et al. Use of hydroxyurea and alpha-interferon in chronic myeloid leukemia during pregnancy: a case report. *Eur J Gynaecol Oncol* 2000;21:89-90.
- ²⁴ Thauvin-Robinet C, Maingueneau C, Robert E, et al. Exposure to hydroxyurea during pregnancy: a case series. *Leukemia* 2001;15:1309-1311.
- ²⁵ Fadilah SA, Ahmad-Zailani H, Soon-Keng C, Norlaila M. Successful treatment of chronic myeloid leukemia during pregnancy with hydroxyurea. *Leukemia* 2002;16:1202-1203.
- ²⁶ Koh LP, Kanagalingam D. Pregnancies in patients with chronic myeloid leukemia in the era of imatinib. *Int J Hematol* 2006;84:459-462.
- ²⁷ Ali R, Ozkalemkas F, Ozkocaman V, et al. Successful pregnancy and delivery in a patient with chronic myelogenous leukemia (CML), and management of CML with leukapheresis during pregnancy: a case report and review of the literature. *Jpn J Clin Oncol* 2004;34:215-217.
- ²⁸ Palani R, Milojkovic D, Apperley JF. Managing pregnancy in chronic myeloid leukaemia. *Ann Hematol* 2015;94 Suppl 2:S167-176.
- ²⁹ Staley EM, Simmons SC, Feldman AZ, et al. Management of chronic myeloid leukemia in the setting of pregnancy: when is leukocytapheresis appropriate? A case report and review of the literature. *Transfusion* 2018;58:456-460.
- ³⁰ James AH, Brancazio LR, Price T. Aspirin and reproductive outcomes. *Obstet Gynecol Surv* 2008;63:49-57.
- ³¹ Deruelle P, Coulon C. The use of low-molecular-weight heparins in pregnancy--how safe are they? *Curr Opin Obstet Gynecol* 2007;19:573-577.
- ³² Russell MA, Carpenter MW, Akhtar MS, et al. Imatinib mesylate and metabolite concentrations in maternal blood, umbilical cord blood, placenta and breast milk. *J Perinatol* 2007;27:241-243.
- ³³ Ali R, Ozkalemkas F, Kimya Y, et al. Imatinib use during pregnancy and breast feeding: a case report and review of the literature. *Arch Gynecol Obstet* 2009;280:169-175.
- ³⁴ Drugs and Lactation Database (LactMed) [Internet]. Bethesda (MD): National Library of Medicine (US); 2006-.
- ³⁵ Chelysheva E, Aleshin S, Polushkina E, et al. Breastfeeding in patients with chronic myeloid leukaemia: Case series with measurements of drug concentrations in maternal milk and literature review. *Mediterr J Hematol Infect Dis* 2018;10:e2018027.
- ³⁶ Abruzzese E, Trawinska MM, Perrotti AP, De Fabritiis P. Tyrosine kinase inhibitors and pregnancy. *Mediterr J Hematol Infect Dis* 2014;6:e2014028.

Note: All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.



NCCN Guidelines Version 2.2022

Chronic Myeloid Leukemia

CRITERIA FOR RESPONSE AND RELAPSE

Response/Relapse	Definition
Complete hematologic response (CHR)¹	• Complete normalization of peripheral blood counts with leukocyte count $<10 \times 10^9/L$ • Platelet count $<450 \times 10^9/L$ • No immature cells, such as myelocytes, promyelocytes, or blasts in peripheral blood • No signs and symptoms of disease with resolution of palpable splenomegaly
Cytogenetic response^{2,3}	• Complete cytogenetic response (CCyR): No Ph-positive metaphases⁴ • Major cytogenetic response (MCyR): 0%–35% Ph-positive metaphases • Partial cytogenetic response (PCyR): 1%–35% Ph-positive metaphases • Minor cytogenetic response: $>35\%$–65% Ph-positive metaphases
Molecular response^{5,6,7}	• Early molecular response (EMR): <i>BCR-ABL1</i> (IS) $\leq 10\%$ at 3 and 6 months • Major molecular response (MMR): <i>BCR-ABL1</i> (IS) $\leq 0.1\%$ or ≥ 3-log reduction in <i>BCR-ABL1</i> transcripts from the standardized baseline, if qPCR (IS) is not available • Deep molecular response (DMR): MR4.0: <i>BCR-ABL1</i> (IS) $\leq 0.01\%$ or MR4.5: <i>BCR-ABL1</i> (IS) $\leq 0.0032\%$
Relapse	• Any sign of loss of hematologic response • Any sign of loss of CCyR or its molecular response correlate defined as an increase in <i>BCR-ABL1</i> transcript to $>1\%$⁴ • 1-log increase in <i>BCR-ABL1</i> transcript levels with loss of MMR⁸

¹ Faderl S, Talpaz M, Estrov Z, Kantarjian HM. Chronic myelogenous leukemia: biology and therapy. *Ann Intern Med* 1999;131:207-219. The American College of Physicians-American Society of Internal Medicine is not responsible for the accuracy of the translation.

² A minimum of 20 metaphases should be examined.

³ O'Brien SG, Guilhot F, Larson RA, et al. Imatinib compared with interferon and low-dose cytarabine for newly diagnosed chronic-phase chronic myeloid leukemia. *N Engl J Med* 2003;348:994-1004.

⁴ CCyR correlates with *BCR-ABL1* (IS) $\leq 1\%$.

⁵ Hughes TP, Kaeda J, Branford S, et al. Frequency of major molecular responses to imatinib or interferon alfa plus cytarabine in newly diagnosed chronic myeloid leukemia. *N Engl J Med* 2003;349:1423-1432.

⁶ Hughes T, Deininger M, Hochhaus A, et al. Monitoring CML patients responding to treatment with tyrosine kinase inhibitors: review and recommendations for harmonizing current methodology for detecting *BCR-ABL* transcripts and kinase domain mutations and for expressing results. *Blood* 2006;108:28-37.

⁷ Cross NC, White HE, Müller MC, Saglio G, Hochhaus A. Standardized definitions of molecular response in chronic myeloid leukemia. *Leukemia* 2012;26:2172-2175.

⁸ The loss of MMR in the presence of a CCyR does not necessarily constitute treatment failure.

Note: All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.



NCCN Guidelines Version 2.2022

Chronic Myeloid Leukemia

MONITORING RESPONSE TO TKI THERAPY AND MUTATIONAL ANALYSIS

Test	Recommendation
Bone marrow cytogenetics ¹	• At diagnosis • Failure to reach response milestones • Any sign of loss of hematologic response • Any sign of loss of CCyR or its molecular response correlate defined as an increase in <i>BCR-ABL1</i> transcript to >1%
qPCR using IS	• At diagnosis • Every 3 months after initiating treatment. After <i>BCR-ABL1</i> (IS) ≤1%² has been achieved, every 3 months for 2 years and every 3–6 months thereafter • If there is a 1-log increase in <i>BCR-ABL1</i> transcript levels with MMR, qPCR should be repeated in 1–3 months
<i>BCR-ABL1</i> kinase domain mutation analysis	• Chronic phase ▸ Failure to reach response milestones ▸ Any sign of loss of hematologic response ▸ Any sign of loss of CCyR or its molecular response correlate defined as an increase in <i>BCR-ABL1</i> transcript to >1% ▸ 1-log increase in <i>BCR-ABL1</i> transcript levels and loss of MMR • Disease progression to accelerated or blast phase³

¹ FISH has been inadequately studied for monitoring response to treatment.

² CCyR correlates with *BCR-ABL1* (IS) ≤1%.

³ Consider myeloid mutation panel to identify *BCR-ABL1*–independent resistance mutations in patients with no *BCR-ABL1* kinase domain mutations.

Note: All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.

**DISCONTINUATION OF TKI THERAPY****General Considerations**

- Discontinuation of TKI therapy appears to be safe in select CML patients.
- Consult with a CML specialist to review the appropriateness for TKI discontinuation and potential risks and benefits of treatment discontinuation, including TKI withdrawal syndrome.
- Clinical studies that have evaluated the safety and efficacy of TKI discontinuation have employed strict eligibility criteria and have mandated more frequent molecular monitoring than typically recommended for patients on TKI therapy.
- Some patients have experienced significant adverse events that are believed to be due to TKI discontinuation.
- Discontinuation of TKI therapy should only be performed in consenting patients after a thorough discussion of the potential risks and benefits.
- Consultation with an NCCN Panel Member or center of expertise is recommended in the following circumstances:
 - Any significant adverse event is believed to be related to treatment discontinuation.
 - There is progression to accelerated or blast phase CML at any time.
 - There is failure to regain MMR after 3 months following treatment reinitiation.
- Outside of a clinical trial, TKI discontinuation should be considered only if ALL of the criteria included in the list below are met.

Criteria for TKI Discontinuation

- Age ≥ 18 years.
- Chronic phase CML. No prior history of accelerated or blast phase CML.
- On approved TKI therapy for at least 3 years.^{1,2}
- Prior evidence of quantifiable *BCR-ABL1* transcript.
- Stable molecular response (MR4; *BCR-ABL1* $\leq 0.01\%$ IS) for ≥ 2 years, as documented on at least 4 tests, performed at least 3 months apart.²
- Access to a reliable qPCR test with a sensitivity of detection of at least MR4.5 (*BCR-ABL1* $\leq 0.0032\%$ IS) and that provides results within 2 weeks.
- Monthly molecular monitoring for the first 6 months following discontinuation, bimonthly during months 7–12, and quarterly thereafter (indefinitely) for patients who remain in MMR (MR3; *BCR-ABL1* $\leq 0.1\%$ IS).
- Prompt resumption of TKI within 4 weeks of a loss of MMR with monthly molecular monitoring until MMR is re-established, then every 3 months thereafter is recommended indefinitely for patients who have reinitiated TKI therapy after a loss of MMR. For those who fail to achieve MMR after 3 months of TKI resumption, *BCR-ABL1* kinase domain mutation testing should be performed, and monthly molecular monitoring should be continued for another 6 months.

¹ The feasibility of TFR following discontinuation of TKIs other than dasatinib, imatinib, or nilotinib has not yet been evaluated in clinical studies. It is reasonable to assume that the likelihood of TFR following discontinuation would be similar irrespective of TKI in patients who have achieved and maintained deep molecular response (MR4.0; $\leq 0.01\%$ *BCR-ABL1* IS) for ≥ 2 years, based on the extrapolation of findings from the studies that have evaluated TFR following discontinuation of imatinib, dasatinib, or nilotinib.

² Data from the EURO-SKI study suggest that MR4.0 (*BCR-ABL1* $\leq 0.01\%$ IS) for ≥ 3 years was the most significant predictor for successful discontinuation of imatinib. Total duration of imatinib therapy for at least 6 years was also predictive of successful discontinuation (Saussele S, et al. Lancet Oncol 2018;19:747-757).

Note: All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.



MANAGEMENT OF TOXICITIES AND DRUG INTERACTIONS

[BOSUTINIB \(CML-G 1 of 9\)](#)

[DASATINIB \(CML-G 2 of 9\)](#)

[IMATINIB \(CML-G 3 of 9\)](#)

[NILOTINIB \(CML-G 4 of 9\)](#)

[OMACETAXINE \(CML-G 5 of 9\)](#)

[PONATINIB \(CML-G 6 of 9\)](#)

[ASCIMINIB \(CML-G 7 of 9\)](#)

[DRUG INTERACTIONS OF TKIs \(CML-G 8 of 9\)](#)

Note: All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.

**MANAGEMENT OF BOSUTINIB TOXICITY¹****Dosing:**

Newly diagnosed CP-CML: The recommended initial dose is 400 mg once daily.

CP-CML, AP-CML or BP-CML with resistance or intolerance to prior therapy: The recommended initial dose is 500 mg once daily.

Dose Adjustments:**Hematologic Toxicities**

- **Absolute neutrophil count (ANC) $<1.0 \times 10^9/L$ or platelets $<50 \times 10^9/L$:** Hold bosutinib until ANC $\geq 1.0 \times 10^9/L$ and platelets $\geq 50 \times 10^9/L$. Resume treatment with bosutinib at the same dose if recovery occurs within 2 weeks. If blood counts remain low for greater than 2 weeks, upon recovery reduce dose by 100 mg and resume treatment. If cytopenia recurs, reduce dose by an additional 100 mg upon recovery and resume treatment. Doses less than 300 mg/day have not been evaluated.
- **Growth factors can be used in combination with bosutinib for patients with persistent neutropenia and thrombocytopenia.**
- **Grade 3–4 anemia:²** Check reticulocyte count, ferritin, iron saturation, B12, folate, and correct nutritional deficiencies if present. Transfusion support should be used if patient is symptomatic.

Non-Hematologic Toxicities

- **Liver transaminases $>5 \times$ institutional upper limit of normal (IULN):** Hold bosutinib until recovery to $\leq 2.5 \times$ IULN and resume dose at 400 mg once daily thereafter. If recovery takes longer than 4 weeks, discontinue bosutinib. If transaminase elevations $\geq 3 \times$ IULN occur concurrently with bilirubin elevations $>2 \times$ IULN and alkaline phosphatase $<2 \times$ IULN (Hy's law case definition), discontinue bosutinib.
- **Diarrhea:** For NCI Common Terminology Criteria for Adverse Events (CTCAE) grade 3–4 diarrhea (increase of ≥ 7 stools/day over baseline/pretreatment), withhold bosutinib until recovery to grade ≤ 1 . Bosutinib may be resumed at 400 mg once daily.
- **For other clinically significant, moderate, or severe non-hematologic toxicity,** withhold bosutinib until the toxicity has resolved, then consider resuming bosutinib at 400 mg once daily. If clinically appropriate, consider re-escalating the dose of bosutinib to 500 mg once daily.

Special Populations

- **In patients with pre-existing mild, moderate, and severe hepatic impairment,** the recommended dose of bosutinib is 200 mg daily. A daily dose of 200 mg in patients with hepatic impairment is predicted to result in an area under the curve (AUC) similar to the AUC seen in patients with normal hepatic function receiving 500 mg daily. However, there are no clinical data for efficacy at the dose of 200 mg once daily in patients with hepatic impairment and CML.
- **Monitoring renal function at baseline and during therapy is recommended with particular attention to those patients with preexisting renal impairment or risk factors for renal dysfunction.** Dose modification (as outlined in the prescribing information) should be considered for patients with baseline and treatment-emergent renal impairment.

Specific Interventions

- **Fluid retention events (ie, pulmonary and/or peripheral edema; pleural and pericardial effusion):** Diuretics, supportive care.
- **GI upset:** Take medication with a meal and large glass of water.
- **Rash:** Topical or systemic steroids, dose reduction, dose interruption, or dose discontinuation.

¹ Please refer to package insert for full prescribing information and monitoring of hematologic or biochemical abnormalities:

<https://www.accessdata.fda.gov/scripts/cder/daf/index.cfm>.

² Although erythropoietin is effective, guidelines from the Centers for Medicare & Medicaid Services (CMS) and the U.S. Food and Drug Administration (FDA) do not support the use of erythropoiesis-stimulating agents (ESAs) in myeloid malignancies.

Note: All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.

**MANAGEMENT OF DASATINIB TOXICITY¹****Dosing:**

- CP-CML: The recommended initial dose is 100 mg once daily.
- AP-CML and BP-CML: The recommended initial dose is 140 mg once daily.

Dose Adjustments:**Hematologic Toxicities**

- Chronic phase ANC $<0.5 \times 10^9/L$ or platelets $<50 \times 10^9/L$: Hold dasatinib until ANC $\geq 1.0 \times 10^9/L$ and platelets $\geq 50 \times 10^9/L$, then resume dasatinib at the starting dose if recovery occurs in ≤ 7 days. If platelets $<25 \times 10^9/L$ or recurrence of ANC $<0.5 \times 10^9/L$ for >7 days, hold drug until ANC $\geq 1.0 \times 10^9/L$ and platelets $\geq 50 \times 10^9/L$, then resume dasatinib at reduced dose of 80 mg once daily for second episode. For third episode, further reduce dose to 50 mg once daily (for newly diagnosed patients) or discontinue dasatinib (for patients with disease that is resistant or intolerant to prior therapy including imatinib).
- Accelerated phase and blast phase, ANC $<0.5 \times 10^9/L$ and/or platelets $<10 \times 10^9/L$: Patients may have cytopenias related to disease. If cytopenia is unrelated to disease, hold dasatinib until ANC $\geq 1.0 \times 10^9/L$ and platelets $\geq 20 \times 10^9/L$, and resume at original starting dose. If recurrence, hold dasatinib until ANC $\geq 1.0 \times 10^9/L$ and platelets $\geq 20 \times 10^9/L$, and resume dasatinib at reduced dose of 100 mg once daily (second episode) or 80 mg once daily (third episode).
- Growth factors can be used in combination with dasatinib for patients with persistent neutropenia and thrombocytopenia.
- Grade 3–4 anemia:² Check reticulocyte count, ferritin, iron saturation, B12, folate, and correct nutritional deficiencies if present. Transfusion support should be used if patient is symptomatic.

Non-Hematologic Toxicities

- If a severe, non-hematologic, adverse reaction develops with dasatinib, treatment must be held until the event has resolved or improved. Thereafter, treatment can be resumed as appropriate at a reduced dose depending on the initial severity of the event.

Rare But Serious Toxicities

- Pulmonary arterial hypertension (PAH): Dasatinib can cause PAH, which may occur any time after initiation, including after more than one year of treatment. PAH may be reversible on discontinuation of dasatinib. Evaluate patients for signs and symptoms of underlying cardiopulmonary disease prior to initiating dasatinib and during treatment. If PAH is confirmed, dasatinib should be permanently discontinued.

Specific Interventions

- Fluid retention events (ie, ascites, edema, pleural and pericardial effusion): Diuretics, supportive care.
- Pleural/pericardial effusion: Diuretics, dose interruption. If patient has significant symptoms, consider short course of steroids (prednisone 20–50 mg/day x 3–4 days, may taper with 20 mg/day x 3–4 days); when resolved, reduce one dose level.
- GI upset: Take medication with a meal and large glass of water.
- Rash: Topical or systemic steroids, dose reduction, dose interruption, or dose discontinuation.

¹ Please refer to package insert for full prescribing information and monitoring of hematologic or biochemical abnormalities:

<https://www.accessdata.fda.gov/scripts/cder/daf/index.cfm>.

² Although erythropoietin is effective, guidelines from CMS and the FDA do not support the use of ESAs in myeloid malignancies.

Note: All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.

**MANAGEMENT OF IMATINIB TOXICITY^{1,3}****Dosing:**

- **CP-CML:** The recommended initial dose is 400 mg once daily.
- **AP-CML and BP-CML:** The recommended initial dose is 600 mg once daily.

Dose Adjustments:**Hematologic Toxicities**

- **Chronic phase ANC $<1.0 \times 10^9/L$, and/or platelets $<50 \times 10^9/L$:** Hold imatinib until ANC $\geq 1.5 \times 10^9/L$ and platelets $\geq 75 \times 10^9/L$, then resume imatinib at the starting dose of 400 mg. If recurrence of ANC $<1.0 \times 10^9/L$ and/or platelets $<50 \times 10^9/L$, hold drug until ANC $\geq 1.5 \times 10^9/L$ and platelets $\geq 75 \times 10^9/L$, then resume imatinib at reduced dose of 300 mg.
- **Accelerated phase and blast phase, ANC $<0.5 \times 10^9/L$ and/or platelets $<10 \times 10^9/L$:** Patients may have cytopenias related to disease. If cytopenia is unrelated to disease, reduce dose to 400 mg. If cytopenia persists for 2 weeks, reduce dose further to 300 mg. If cytopenia persists for 4 weeks, stop imatinib until ANC $\geq 1.0 \times 10^9/L$ and platelet count $\geq 20 \times 10^9/L$ and then resume treatment at 300 mg.
- **Growth factors** can be used in combination with imatinib for patients with persistent neutropenia.⁴
- **Grade 3–4 anemia:**² Check reticulocyte count, ferritin, iron saturation, B12, folate, and correct nutritional deficiencies if present. Transfusion support should be used if patient is symptomatic.

Non-Hematologic Toxicities

- **Bilirubin $>3 \times IULN$ or liver transaminases $>5 \times IULN$:** Hold imatinib until bilirubin $<1.5 \times IULN$ and transaminase levels $<2.5 \times IULN$. Resume imatinib at a reduced daily dose (400–300 mg, 600–400 mg, or 800–600 mg).
- **Severe hepatotoxicity or severe fluid retention:** Hold imatinib until the event has resolved. Treatment can be resumed as appropriate depending on the severity of the event.
- **Patients with moderate renal impairment (creatinine clearance [CrCl] = 20–39 mL/min)** should receive a 50% decrease in the recommended starting dose and future doses can be increased as tolerated. Doses greater than 600 mg are not recommended in patients with mild renal impairment (CrCl = 40–59 mL/min). For patients with moderate renal impairment, doses greater than 400 mg are not recommended. Imatinib should be used with caution in patients with severe renal impairment.

Specific Interventions

- **Fluid retention (ie, pleural effusion, pericardial effusion, edema, ascites):** Diuretics, supportive care, dose reduction, interruption, or discontinuation. Consider echocardiogram to check left ventricular ejection fraction (LVEF).
- **GI upset:** Take medication with a meal and large glass of water.
- **Muscle cramps:** Calcium supplement, tonic water.
- **Rash:** Topical or systemic steroids, dose reduction, dose interruption, or dose discontinuation.

¹ Please refer to package insert for full prescribing information and monitoring of hematologic or biochemical abnormalities:

<https://www.accessdata.fda.gov/scripts/cder/daf/index.cfm>.

² Although erythropoietin is effective, guidelines from CMS and the FDA do not support the use of ESAs in myeloid malignancies.

³ Many toxicities are self-limiting; consider re-escalating dose at a later time.

⁴ Quintas-Cardama A, Kantarjian H, O'Brien S, et al. Granulocyte-colony-stimulating factor (filgrastim) may overcome imatinib-induced neutropenia in patients with chronic-phase chronic myelogenous leukemia. Cancer 2004;100:2592-2597.

Note: All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.

**MANAGEMENT OF NILOTINIB TOXICITY¹**

- Nilotinib prolongs the QT interval. Prior to administration of nilotinib and periodically, monitor for hypokalemia or hypomagnesemia and correct deficiencies. Electrocardiograms (ECGs) should be obtained to monitor the QTc at baseline, 7 days after initiation, and periodically thereafter, as well as following any dose adjustments.
- Sudden deaths have been reported in patients receiving nilotinib.
- Avoid use of concomitant drugs known to prolong the QT interval and strong CYP3A4 inhibitors.
- Patients should avoid food 2 hours before and 1 hour after taking dose.

Dosing:

- Newly diagnosed CP-CML: The recommended initial dose is 300 mg twice daily.
- Resistant or intolerant CP-CML and AP-CML: The recommended initial dose is 400 mg twice daily.
- BP-CML: The recommended initial dose is 400 mg twice daily.

Dose Adjustments:**QT Interval Prolongation**

- ECGs with a QTc >480 msec: Hold nilotinib. If serum potassium and magnesium levels are below lower limit of normal, correct with supplements to within normal limits. Review concomitant medication usage. Resume within 2 weeks at prior dose if QTcF is <450 msec and within 20 msec of baseline. If QTcF is between 450 and 480 msec after 2 weeks, resume at reduced dose (400 mg once daily). Following dose reduction, if QTcF returns to >480 msec, nilotinib should be discontinued. ECG should be obtained 7 days after any dose adjustment to monitor QTc.

Hematologic Toxicities

- Chronic or accelerated phase, ANC <1.0 x 10⁹/L, and/or platelets <50 x 10⁹/L: Hold nilotinib and monitor blood counts. Resume within 2 weeks at prior dose if ANC >1.0 x 10⁹/L and platelets >50 x 10⁹/L. If blood counts remain low for >2 weeks, reduce dose to 400 mg once daily.
- Growth factors can be used in combination with nilotinib for patients with persistent neutropenia and thrombocytopenia.
- Grade 3–4 anemia:² Check reticulocyte count, ferritin, iron saturation, B12, folate, and correct nutritional deficiencies if present. Transfusion support should be used if patient is symptomatic.

¹ Please refer to package insert for full prescribing information and monitoring of hematologic or biochemical abnormalities:

<https://www.accessdata.fda.gov/scripts/cder/daf/index.cfm>.

² Although erythropoietin is effective, guidelines from CMS and the FDA do not support the use of ESAs in myeloid malignancies.

Non-Hematologic Toxicities

- Elevated serum lipase, amylase, bilirubin, or hepatic transaminases grade ≥3: Hold nilotinib and monitor serum levels. Resume nilotinib at 400 mg once daily if serum levels return to grade ≤1.
- Hepatic Impairment: Consider alternate therapies. See prescribing information for dose adjustments related to hepatic impairment.
- Assess glucose levels before initiating treatment and monitor treatment as clinically indicated.

Rare But Serious Toxicities

- Peripheral arterial occlusive disease (PAOD): Nilotinib is associated with an increased risk of vascular adverse events, including PAOD, and should be used with caution in patients with cardiovascular risk factors or a history of PAOD. Evaluate patients for a history of PAOD and for vascular risk factors prior to initiating nilotinib and during treatment. If PAOD is confirmed, nilotinib should be permanently discontinued.

Specific Interventions

- Rash: Topical or systemic steroids, dose reduction, dose interruption, or dose discontinuation.

Note: All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.



MANAGEMENT OF OMACETAXINE TOXICITY¹

Dosing:

- **Resistant or intolerant CP-CML and AP-CML:**
 - ▶ **Induction dose:** The recommended initial dose is 1.25 mg/m² twice daily for 14 consecutive days of a 28-day cycle.
 - ▶ **Maintenance dose:** 1.25 mg/m² twice daily for 7 consecutive days of a 28-day cycle.

Dose Adjustments:

Hematologic Toxicities

- CBCs should be performed weekly during induction and initial maintenance cycles. After initial maintenance cycles, monitor CBCs every 2 weeks or as clinically indicated. ANC <0.5 x 10⁹/L or platelet count <50 x 10⁹/L: Delay starting the next cycle until ANC ≥1.0 x 10⁹/L and platelet count ≥50 x 10⁹/L and reduce the number of dosing days by 2 days for the next cycle.

Non-Hematologic Toxicities

- **Grade 3 or 4 hyperglycemia:** Monitor blood glucose levels frequently, especially in patients with diabetes or risk factors for diabetes. Avoid omacetaxine in patients with poorly controlled diabetes mellitus (DM) until good glycemic control has been established.
- **Manage other clinically significant non-hematologic toxicity symptomatically.** Interrupt and/or delay omacetaxine until toxicity is resolved.

¹ Please refer to package insert for full prescribing information and monitoring of hematologic or biochemical abnormalities:
<https://www.accessdata.fda.gov/scripts/cder/daf/index.cfm>.

Note: All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.

**MANAGEMENT OF PONATINIB TOXICITY¹**

- Arterial occlusive events (AOEs; including fatal myocardial infarction and stroke) and venous thromboembolic events (VTEs), have occurred in patients treated with ponatinib. Monitor for evidence of thromboembolism and vascular occlusion. Interrupt or stop ponatinib immediately for vascular occlusion.
- Heart failure has occurred in patients treated with ponatinib. Monitor cardiac function. Interrupt or stop ponatinib for new or worsening heart failure.
- Hepatotoxicity: Hepatotoxicity, liver failure, and death have occurred in patients treated with ponatinib. Monitor hepatic function prior to and during treatment. Interrupt ponatinib if hepatotoxicity is suspected.
- Cardiovascular risk: Identify and control traditional risk factors for atherosclerosis (eg, DM, hypertension, hyperlipidemia, smoking, estrogen use) before starting ponatinib. Patients with cardiovascular risk factors should be referred to a cardiologist. Consider the use of low-dose aspirin if there is no contraindication.
- Ponatinib is also associated with grade ≥3 skin rash and pancreatitis leading to dose modifications (dose delays or dose reductions).

Dosing:

- CP-CML: The recommended initial dose is 45 mg once daily with a reduction to 15 mg once daily upon achievement of *BCR-ABL1* (IS) ≤1%. In patients with a loss of response, the dose can be re-escalated to a previously tolerated dose of 30 mg or 45 mg once daily and continued at the re-escalated dose until loss of response or unacceptable toxicity.
- AP-CML or BP-CML: The recommended initial dose is 45 mg once daily. Consider reducing the dose for patients with AP-CML who have achieved a MCyR. Consider discontinuing ponatinib if response has not occurred by 3 months.

Dose Adjustments:**Hematologic Toxicities**

- ANC <1.0 x 10⁹/L or platelets <50 x 10⁹/L
 - ▶ First occurrence: Hold ponatinib until ANC ≥1.5 x 10⁹/L and platelets ≥75 x 10⁹/L and resume at initial dose of 45 mg.
 - ▶ Second occurrence: Hold ponatinib until ANC ≥1.5 x 10⁹/L and platelets ≥75 x 10⁹/L and resume at 30 mg.
 - ▶ Third occurrence: Hold ponatinib until ANC ≥1.5 x 10⁹/L and platelets ≥75 x 10⁹/L and resume at 15 mg.
- Growth factors can be used in combination with ponatinib for patients with persistent neutropenia and thrombocytopenia.
- Grade 3–4 anemia:² Check reticulocyte count, ferritin, iron saturation, B12, folate, and correct nutritional deficiencies if present. Transfusion support should be used if patient is symptomatic.

Non-Hematologic Toxicities

- Liver transaminase >3 x ULN (grade ≥2): Monitor hepatic function. Hold drug until serum levels are <3 x IULN. Resume at lower dose after recovery (30 mg if patient is receiving 45 mg; 15 mg if patient is receiving 30 mg). Discontinue ponatinib if patient is receiving 15 mg.
- AST or ALT ≥3 x ULN concurrent with bilirubin >2 x ULN and alkaline phosphatase <2 x ULN: Discontinue ponatinib.

- Serum lipase elevation, grade 1 or 2 (asymptomatic): Consider dose interruption or reduction. Serum lipase elevation, grade 3 or 4 (>2 x IULN) (asymptomatic) or asymptomatic radiologic pancreatitis: Hold drug until serum levels are <1.5 x ULN. Resume at lower dose after recovery (30 mg if patient is receiving 45 mg; 15 mg if patient is receiving 30 mg). Discontinue ponatinib if patient is receiving 15 mg.
- Pancreatitis (symptomatic), grade 3: Hold drug until serum lipase levels are ≤grade 1. Resume at lower dose after recovery (30 mg if patient is receiving 45 mg; 15 mg if patient is receiving 30 mg). Discontinue ponatinib if patient is receiving 15 mg. Grade 4: Discontinue ponatinib.

Rare But Serious Toxicities

- Hemorrhage: Hemorrhagic events were reported in clinical trials. Cerebral and gastrointestinal hemorrhage were the most commonly reported serious bleeding events. Serious hemorrhage should be managed with dose interruption.
- Cardiac arrhythmias: Advise patients to report signs and symptoms suggestive of alterations in heart rate (fainting, dizziness, chest pain, or palpitations).
- Tumor lysis syndrome: Ensure adequate hydration and correct high uric acid levels prior to initiating therapy with ponatinib in patients with advanced phase CML.

Specific Interventions

- Fluid retention events (ie, edema, ascites, pleural and pericardial effusion) are managed with dose interruption, dose reduction, or discontinuation of ponatinib as clinically indicated.
- Hypertension: Monitor and manage blood pressure elevations.
- Rash: Topical or systemic steroids, dose reduction, dose interruption, or dose discontinuation.

¹ Please refer to package insert for full prescribing information and monitoring of hematologic or biochemical abnormalities: <https://www.accessdata.fda.gov/scripts/cder/daf/index.cfm>.

² Although erythropoietin is effective, guidelines from CMS and the FDA do not support the use of ESAs in myeloid malignancies.

Note: All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.

**MANAGEMENT OF ASCIMINIB TOXICITY¹****Dosing:**

- Asciminib should be taken orally without food. Patients should avoid food 2 hours before and 1 hour after taking asciminib.
- CP-CML (previously treated with ≥ 2 TKIs): The recommended initial dose is 80 mg orally once daily or 40 mg twice daily (at approximately 12-hour intervals).
- CP-CML with T315I mutation: The recommended initial dose is 200 mg orally twice daily (at approximately 12-hour intervals).

Dose Adjustments:**Hematologic Toxicities**

- ANC $< 1.0 \times 10^9/L$, and/or platelets $< 50 \times 10^9/L$: Hold asciminib until ANC $\geq 1.0 \times 10^9/L$ and/or platelets $\geq 50 \times 10^9/L$. Resume asciminib at starting dose (if resolved within 2 weeks) or at reduced dose (if resolved after 2 weeks).
- For recurrent severe thrombocytopenia and/or neutropenia, withhold asciminib until resolved to ANC $\geq 1.0 \times 10^9/L$ and platelets $\geq 50 \times 10^9/L$, then resume at reduced dose.

Non-Hematologic Toxicities

- Elevated serum amylase and/or lipase ($> 2.0 \times \text{ULN}$): Hold asciminib and resume asciminib at reduced dose, after serum levels return to $< 1.5 \times \text{ULN}$. If events reoccur at reduced dose or if serum levels do not return to $< 1.5 \times \text{ULN}$, permanently discontinue asciminib. Perform diagnostic tests to exclude pancreatitis.
- Hypertension: Monitor blood pressure and manage hypertension as clinically indicated. Interrupt, dose reduce, or permanently discontinue asciminib if hypertension is not medically controlled.
- Hypersensitivity: May cause hypersensitivity reactions. Monitor patients for signs and symptoms and initiate appropriate treatment as clinically indicated.
 - Grade ≥ 3 : Hold asciminib until recovery to grade ≤ 1 and resume asciminib at reduced dose. If not resolved, permanently discontinue asciminib.
- Cardiovascular Toxicity: Monitor patients with history of cardiovascular risk factors for cardiovascular signs and symptoms. Initiate appropriate treatment as clinically indicated.
 - Grade ≥ 3 : Hold asciminib until recovery to grade ≤ 1 and resume asciminib at reduced dose. If not resolved, permanently discontinue asciminib.

Drug Interactions:

- Strong CYP3A4 inhibitors: Closely monitor for adverse reactions during concomitant use of asciminib 200 mg twice daily dosing.
- Itraconazole oral solution containing hydroxypropyl- β -cyclodextrin: Avoid concomitant use of asciminib at all recommended doses.
- CYP3A4 substrates: Closely monitor for adverse reactions during concomitant use of asciminib at 80 mg total daily dose. Avoid concomitant use of asciminib 200 mg twice daily dosing.
- CYP2C9 substrates: Avoid concomitant use of asciminib at all recommended doses. If concomitant use is unavoidable, reduce CYP2C9 substrate dosage for use with asciminib 80 mg total daily dose or consider alternative therapy with non-CYP2C9 substrate for use with asciminib 200 mg twice daily dosing.
- P-glycoprotein substrates: Closely monitor for adverse reactions during concomitant use of asciminib at all recommended doses.

¹ Please refer to package insert for full prescribing information and monitoring of hematologic or biochemical abnormalities: <https://www.accessdata.fda.gov/scripts/cder/daf/index.cfm>.

Note: All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.



NCCN Guidelines Version 2.2022

Chronic Myeloid Leukemia

DRUG INTERACTIONS OF TKIs WITH MOST COMMONLY USED DRUGS AND SUPPLEMENTS^{1,5}

Drug interactions with TKIs are not uncommon. It is always important to take a detailed medication history (including herbal supplements) at every visit.

Drug Class/ Medications	Change in TKI Level				
	Bosutinib	Dasatinib	Imatinib	Nilotinib	Ponatinib
Proton Pump Inhibitors (PPIs) • Lansoprazole • Rabeprazole • Esomeprazole • Omeprazole • Pantoprazole	Decrease in exposure	Decrease in exposure	No major interaction	Decrease in exposure	Minor decrease in exposure; Monitor
Histamine 2 Receptor Antagonists (H2RAs) • Famotidine • Ranitidine • Nizatidine	Decrease in exposure; AVOID; If absolutely necessary consider once-daily H2RA ≥2 hours after taking bosutinib	Decrease in exposure; AVOID; If absolutely necessary consider once-daily H2RA ≥2 hours after taking dasatinib	No major interaction	Decrease in exposure; AVOID; If absolutely necessary consider once-daily H2RA ≥2 hours after or ≥10 hours before taking nilotinib	No major interaction
Antacids	Decrease in exposure if concomitant; Use antacids at least 2 hours before or at least 2 hours after taking bosutinib	Decrease in exposure if concomitant; Use antacids at least 2 hours before or at least 2 hours after taking dasatinib	No major interaction	Decrease in exposure if concomitant; Use antacids at least 2 hours before or at least 2 hours after taking nilotinib	No major interaction
Antidepressants • Fluoxetine • Bupropion • Citalopram	Minor increase in exposure; Monitor QTc monitoring	Minor increase in exposure; Monitor QTc monitoring	Minor increase in exposure; Monitor QTc monitoring	AVOID if possible due to cumulative QTc prolongation risk	Minor increase in exposure; Monitor QTc monitoring
Cardiovascular Medications • Amiodarone • Diltiazem • Verapamil	Increase in exposure and arrhythmia risk; Strongly consider alternative cardiac medication or TKI dose adjustment	Increase in exposure and arrhythmia risk; Strongly consider alternative cardiac medication or TKI dose adjustment	Increase in exposure; Strongly consider alternative cardiac medication or TKI dose adjustment	Increase in exposure and arrhythmia risk; AVOID	Increase in exposure; Strongly consider alternative cardiac medication or TKI dose adjustment

[Continued on next page](#)

Note: All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.



NCCN Guidelines Version 2.2022

Chronic Myeloid Leukemia

DRUG INTERACTIONS OF TKIs WITH MOST COMMONLY USED DRUGS AND SUPPLEMENTS^{1,5}

Drug interactions with TKIs are not uncommon. It is always important to take a detailed medication history (including herbal supplements) at every visit.

Drug Class/ Medications	Change in TKI Level				
	Bosutinib	Dasatinib	Imatinib	Nilotinib	Ponatinib
Anti-infectives • Azole Antifungals ▶ Fluconazole ≥200 mg ▶ Voriconazole ▶ Itraconazole ▶ Posaconazole ▶ Isavuconazole • Clarithromycin • Telithromycin • Ritonavir	Increase in exposure; Strongly consider alternative anti-infective or TKI dose adjustment	Increase in exposure; Strongly consider alternative anti-infective or TKI dose adjustment	Increase in exposure; Strongly consider alternative anti-infective or TKI dose adjustment	Increase in exposure; Strongly consider alternative anti-infective or TKI dose adjustment	Increase in exposure; Strongly consider alternative anti-infective or TKI dose adjustment
Anti-infectives • Fluoroquinolones ▶ Levofloxacin ▶ Moxifloxacin ▶ Ciprofloxacin	QTc monitoring	QTc monitoring	No major interaction	Use with caution	No major interaction
Herbal Supplements^{6,7} • Curcumin (Turmeric) • Ginkgo Biloba • Green Tea Extract	Increase in exposure; Strongly consider supplement discontinuation	Increase in exposure; Strongly consider supplement discontinuation	Increase in exposure; Strongly consider supplement discontinuation	Increase in exposure; Strongly consider supplement discontinuation	Increase in exposure; Strongly consider supplement discontinuation
Herbal Supplements^{6,7} • St. John's Wort	Decrease in exposure; AVOID	Decrease in exposure; AVOID	Decrease in exposure; AVOID	Decrease in exposure; AVOID	Decrease in exposure; AVOID

¹ Please refer to package insert for full prescribing information and drug interactions: <https://www.accessdata.fda.gov/scripts/cder/daf/index.cfm>.

⁵ van Leeuwen RW, van Gelder T, Mathijssen RH, et al. Drug-drug interactions with tyrosine-kinase inhibitors: a clinical perspective. Lancet Oncol 2014;15:e315-e326.

⁶ Zhang W, Lim LY. Effects of spice constituents on P-glycoprotein-mediated transport and CYP3A4-mediated metabolism in vitro. Drug Metab Dispos 2008;36:1283-1290.

⁷ Scott GN, Elmer GW. Update on natural product-drug interactions. Am J Health Syst Pharm 2002;59:339-347.

Note: All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.



NCCN Categories of Evidence and Consensus

Category 1	Based upon high-level evidence, there is uniform NCCN consensus that the intervention is appropriate.
Category 2A	Based upon lower-level evidence, there is uniform NCCN consensus that the intervention is appropriate.
Category 2B	Based upon lower-level evidence, there is NCCN consensus that the intervention is appropriate.
Category 3	Based upon any level of evidence, there is major NCCN disagreement that the intervention is appropriate.

All recommendations are category 2A unless otherwise indicated.

NCCN Categories of Preference

Preferred intervention	Interventions that are based on superior efficacy, safety, and evidence; and, when appropriate, affordability.
Other recommended intervention	Other interventions that may be somewhat less efficacious, more toxic, or based on less mature data; or significantly less affordable for similar outcomes.
Useful in certain circumstances	Other interventions that may be used for selected patient populations (defined with recommendation).

All recommendations are considered appropriate.



NCCN Guidelines Version 2.2022


Chronic Myeloid Leukemia

Discussion

This discussion corresponds to the NCCN Guidelines for Chronic Myeloid Leukemia. Last updated: January 13, 2021.

Table of Contents

Overview	MS-2
Literature Search Criteria and Guidelines Update Methodology	MS-2
Diagnosis and Workup	MS-3
Additional Evaluation	MS-5
Management of Chronic Phase CML	MS-6
Management of Advanced Phase CML	MS-18
Emerging Treatment Options	MS-21
Management of CML During Pregnancy and Breastfeeding	MS-21
Specific Considerations for Children with CML	MS-23
References	MS-38





NCCN Guidelines Version 2.2022

Chronic Myeloid Leukemia

Overview

Chronic myeloid leukemia (CML) accounts for 15% of adult leukemias. The median age of disease onset is 67 years; however, CML occurs in all age groups (SEER statistics). In 2020, an estimated 8,450 people will be diagnosed with CML in the United States, and 1,130 people will die from the disease.¹

CML is defined by the presence of Philadelphia chromosome (Ph) in a patient with a myeloproliferative neoplasm (MPN). Ph results from a reciprocal translocation between chromosomes 9 and 22 [t(9;22)] that gives rise to a *BCR-ABL1* fusion gene.² In the majority of patients, the chromosomal break points are located in intron 13 or 14 of the *BCR* gene on chromosome 22 (major break point cluster region; *M-BCR*); in the *ABL1* gene they are located between the two alternative *ABL1* exons 1b and 1a, or between *ABL1* exons 1 and 2.^{3,4} Irrespective of the precise *ABL1* breakpoint, splicing almost invariably fuses *ABL1* exon 2 with *BCR* exons 13 or 14, resulting in e13a2 and e14a2 transcripts that code for a protein, p210, with deregulated tyrosine kinase activity, which causes CML. Unusual *BCR-ABL1* transcripts, e1a2 encoding for p190 (involving the minor break point cluster region; *m-BCR*), or e19a2 encoding for p230 (involving the micro break point cluster region; *μ-BCR*) are found infrequently.^{3,4} p190 is usually produced in the setting of Ph-positive acute lymphoblastic leukemia (ALL), and p230 is associated with enhanced neutrophil differentiation. Atypical *BCR-ABL1* transcripts (eg, e13a3, e14a3, e6a2) have also been detected in about 1% to 2% of patients with CML. The proportion of different *BCR-ABL1* transcripts and the impact of *BCR-ABL1* transcript type on response to TKI therapy are discussed on MS-3.

CML occurs in three different phases (chronic, accelerated, and blast phase) and is usually diagnosed in the chronic phase in the developed world. Untreated chronic phase CML (CP-CML) will eventually progress to

accelerated phase CML (AP-CML) or blast phase CML (BP-CML) in 3 to 5 years on average.⁵ Progression to AP-CML and BP-CML bridges a continuum of clinical features (ie, fever, bone pain, spleen size), cytogenetic changes, and blast count. Gene expression profiling has shown a close correlation of gene expression between AP-CML and BP-CML indicating that the bulk of the genetic changes in progression occur in the transition from CP-CML to AP-CML.⁶ The activation of beta-catenin signaling pathway in CML granulocyte-macrophage progenitors (which enhances the self-renewal activity and leukemic potential of these cells) may be a key pathobiologic event in the evolution to BP-CML.⁷

The NCCN Guidelines for CML discuss the clinical management of CML in all three phases (chronic, accelerated, or blast phase). Evaluation for diseases other than CML as outlined in the NCCN Guidelines for MPN is recommended for all patients with *BCR-ABL1*-negative MPN.

Literature Search Criteria and Guidelines Update Methodology

Prior to the update of this version of the NCCN Guidelines® for Chronic Myeloid Leukemia, an electronic search of the PubMed database was performed to obtain key literature in Chronic Myeloid Leukemia since the last guideline update using the following search terms: chronic myeloid leukemia or chronic myelogenous leukemia. The PubMed database was chosen as it remains the most widely used resource for medical literature and indexes peer-reviewed biomedical literature.⁸

The search results were narrowed by selecting studies in humans published in English. Results were confined to the following article types: Clinical Trial, Phase II; Clinical Trial, Phase III; Clinical Trial, Phase IV; Guideline; Randomized Controlled Trial; Meta-Analysis; Systematic Reviews; and Validation Studies.



NCCN Guidelines Version 2.2022

Chronic Myeloid Leukemia

The data from key PubMed articles selected by the panel for review during the Guidelines update meeting as well as articles from additional sources deemed as relevant to these Guidelines have been included in this version of the Discussion section. Recommendations for which high-level evidence is lacking are based on the panel's review of lower-level evidence and expert opinion.

The complete details of the Development and Update of the NCCN Guidelines are available at www.NCCN.org.

Diagnosis and Workup

Initial evaluation should consist of a history and physical exam, including palpation of spleen, complete blood count (CBC) with differential, chemistry profile, and hepatitis B panel. Bone marrow aspirate and biopsy for morphologic and cytogenetic evaluation and quantitative reverse transcriptase polymerase chain reaction (RT-PCR) to establish the presence of quantifiable *BCR-ABL1* mRNA transcripts at baseline are recommended to confirm the diagnosis of CML ([CML-1](#)).

Bone marrow cytogenetics should be done at initial workup to detect additional chromosomal abnormalities in Ph-positive cells (ACA/Ph⁺), also known as clonal cytogenetic evolution.⁹ If bone marrow evaluation is not feasible, fluorescence *in situ* hybridization (FISH) on a peripheral blood specimen with dual probes for *BCR* and *ABL1* genes is an acceptable method to confirm the diagnosis of CML. Interphase FISH is performed on peripheral blood but can be associated with a false-positive rate of 1% to 5% depending on the specific probe used in the assay.¹⁰ Hypermetaphase FISH is more sensitive and can analyze up to 500 metaphases at a time, but it is applicable only to dividing cells in the bone marrow.¹¹ Double-fusion FISH is associated with low false-positive rates and can detect all variant translocations of the Ph-chromosome.¹²

Quantitative RT-PCR (qPCR) should be done at initial workup to establish the presence of quantifiable *BCR-ABL1* mRNA transcripts. qPCR, usually done on peripheral blood, is the most sensitive assay available for the measurement of *BCR-ABL1* mRNA and it can detect one CML cell in a background of $\geq 100,000$ normal cells. qPCR results can be expressed in various ways, for instance as the ratio of *BCR-ABL1* transcript numbers to the number of control gene transcripts.¹³ An International Scale (IS) has been established to standardize molecular monitoring with qPCR across different laboratories with the use of one of three control genes (*BCR*, *ABL1*, or *GUSB*) and a qPCR assay with a sensitivity of at least 4-log reduction from the standardized baseline.¹⁴ In recent years, IS has become the gold standard of expressing qPCR values. More details on monitoring with qPCR using IS are provided on MS-9. Qualitative RT-PCR for the detection of atypical *BCR-ABL1* transcripts should be considered if there is discordance between FISH and qPCR results. See the section on *BCR-ABL1* Transcript Variants in CML below.

BCR-ABL1 transcripts in the peripheral blood at very low levels (1–10 out of 10^8 peripheral blood leukocytes) can be detected in approximately 30% of normal individuals, and the incidence of this increases with age. The risk of developing CML for these individuals is extremely low, and neither continued monitoring nor therapy are indicated.^{15,16}

BCR-ABL1 Transcript Variants in CML

In an international retrospective analysis of a large cohort with new diagnosed CML (more than 45,000 patients), e13a2 and e14a2 transcripts (both encoding for p210) were identified in 38% and 62% of patients, respectively; e13a2 was more frequent in males and the proportion decreased with age in both sexes.¹⁷ Unusual or atypical transcripts were identified in about 2% of patients (e1a2, e19a2, e13a3, and e14a3 were the most frequently identified transcripts).¹⁷ The



NCCN Guidelines Version 2.2022

Chronic Myeloid Leukemia

incidence of these atypical transcripts was higher in females and the proportion decreased with age in both genders.

The presence of e14a2 at baseline was associated with higher molecular response rates to imatinib.¹⁸⁻²¹ While some studies have demonstrated a trend towards better survival outcomes with e14a2 transcript,^{19,20} in other studies the type of transcript did not have any significant impact on long-term survival outcomes.^{18,22} There are very limited data regarding the impact of these transcripts on response to second-generation tyrosine kinase inhibitor (TKI) therapy.¹⁹ In the study that included 213 patients treated with dasatinib or nilotinib, among patients with e13a2 transcripts, cytogenetic and molecular response rates were higher in patients receiving dasatinib or nilotinib compared to those treated with imatinib.¹⁹ These findings suggest that initial treatment with second-generation TKIs might be beneficial for patients with e13a2 transcripts, although this needs to be confirmed in a prospective study.

The presence of e1a2 transcript (encoding for p190) is associated with higher risk of disease progression and inferior cytogenetic and molecular responses to TKI therapy.²³⁻²⁷ In multivariate analysis, e1a2 transcript was also identified as an independent predictor of inferior survival outcomes.²⁵ It is important to be aware that these data refer to the presence of dominant e1a2 transcript, not to the presence of low-level e1a2 transcripts in patients with dominant e13a2 or e14a2 transcripts. The presence of e19a2 transcript (encoding for p230) is associated with lower rates of cytogenetic and molecular response to TKIs and inferior survival outcomes, despite previous reports of an indolent disease course in the pre-TKI era.²⁶⁻²⁸ Referral to centers with expertise in the management of CML is recommended.

Qualitative RT-PCR, nested RT-PCR, or Sanger sequencing are useful for the identification of atypical *BCR-ABL1* transcripts.^{29,30} qPCR using log-reduction from standardized baseline can be used to monitor e1a2

transcripts, and monitoring e19a2 transcripts is usually performed using qualitative RT-PCR or nested RT-PCR. However, there are no standardized qPCR assays for monitoring molecular response to TKI therapy in patients with atypical *BCR-ABL1* transcripts.^{31,32} The utility of multiplex PCR assays and patient-specific genomic DNA quantitative PCR assay for monitoring atypical *BCR-ABL1* transcripts has been demonstrated in some reports.³³⁻³⁷

Clonal Cytogenetic Evolution

The prognostic significance of ACA/Ph⁺ is related to the specific chromosomal abnormality and other features of accelerated phase.³⁸⁻⁴² The presence of “major route” ACA/Ph⁺ (trisomy 8, isochromosome 17q, second Ph, and trisomy 19) at diagnosis may have a negative prognostic impact on survival and disease progression to accelerated or blast phase.⁴³⁻⁴⁵ However, in a more recent analysis that evaluated the outcomes of patients with CP-CML (with or without ACA) treated with TKI therapy in prospective studies, the presence of ACA/Ph⁺ at the time of diagnosis was not associated with worse prognosis.⁴⁶ Survival outcomes were not statistically significant different among patients with ACA/Ph⁺ based on TKI therapy (imatinib vs. second-generation TKIs) or imatinib dose (400 mg vs. 800 mg). It remains uncertain if second-generation TKIs or high-dose imatinib would be more beneficial for patients with ACA/Ph⁺. Patients with ACA/Ph⁺ at diagnosis should be watched carefully for evidence of therapy failure.

Clonal cytogenetic evolution in Ph-negative cells has also been reported in a small subset of patients treated with TKI therapy.⁴⁷⁻⁵⁸ The most common abnormalities include trisomy 8 and loss of Y chromosome. Previous work suggested that the overall prognosis of Ph-negative CML with clonal evolution is good and is dependent on response to imatinib therapy.⁵¹ Recently, however, the presence of chromosome abnormalities other than loss of Y chromosome has been associated with decreased survival in



NCCN Guidelines Version 2.2022

Chronic Myeloid Leukemia

patients with CP-CML treated with various TKIs, suggesting that closer follow-up is indicated.⁵⁹ Progression to myelodysplastic syndromes (MDS) and acute myeloid leukemia (AML) have been reported in patients with monosomy 7 (del 7q).⁶⁰⁻⁶²

Role of NGS

NGS allows for the detection of low-level BCR-ABL1 kinase domain mutations as well as resistance mutations in genes other than *BCR-ABL1* that may confer resistance to TKIs or portend disease progression.⁶³⁻⁶⁶ In a recent, prospective, multicenter study (NEXT-in-CML) that assessed the feasibility of NGS in detection of low-level mutations in 236 consecutive patients with CML and inadequate response to TKI therapy, NGS was more effective than conventional Sanger sequencing in the detection of low-level mutations.⁶⁶ Prospective monitoring of mutation kinetics demonstrated that TKI-resistant low-level mutations are invariably selected if the patients are not switched to another TKI or if they are switched to an inappropriate TKI or TKI dose.⁶⁶ NGS with myeloid mutation panel should be considered for patients with no identifiable *BCR-ABL1* mutations.

Additional Evaluation

Chronic Phase CML (CML-1)

Sokal and Hasford (Euro) scoring systems have been used for the risk stratification of patients into three risk groups (low, intermediate, and high) in clinical trials evaluating TKIs (CML-A).^{67,68} The Sokal score is based on the patient's age, spleen size on clinical exam, platelet count, and percentage of blasts in the peripheral blood.⁶⁷ The Euro score includes eosinophils and basophils in the peripheral blood in addition to the same clinical variables used in the Sokal score.⁶⁸

The European Treatment and Outcome Study long-term survival (ELTS) score is based on the same variables as the Sokal score and provides the

most useful predictor of CML-related death in patients treated with first-line imatinib.⁶⁹ The ELTS score has been validated in a cohort of 1120 patients with CP-CML treated with imatinib in six clinical trials. Higher age, higher peripheral blasts, bigger spleen, and low platelet counts were significantly associated with increased probabilities of dying of CML. Patients in the intermediate- and the high-risk group had significantly higher probabilities of dying of CML than those in the low-risk group, and the probabilities were also significantly different between the intermediate- and high-risk groups. Unlike other scoring systems, the ELTS score is focused on CML-specific overall survival (OS). This is important, as many patients with CML die from non-CML causes, reflecting the efficacy of TKI therapy.

Determination of risk score using either the Sokal or Euro or ELTS scoring systems prior to initiation of TKI therapy is recommended for patients diagnosed with CP-CML.⁶⁷⁻⁶⁹

Advanced Phase CML (CML-1)

The modified MD Anderson Cancer Center criteria for AP-CML (15% and 29% peripheral blood or bone marrow myeloblasts; ≥30% or more of peripheral blood myeloblasts and promyelocytes; ≥20% of peripheral blood or bone marrow basophils; platelet count ≤100 × 10⁹/L unrelated to therapy; and clonal cytogenetic evolution in Ph+ cells) are used in many clinical trials that have evaluated the efficacy of TKIs (CML-B).⁷⁰ The revised 2017 WHO diagnostic criteria for AP-CML include a "provisional" response to TKI criteria in addition to hematologic and cytogenetic criteria.⁷¹ These diagnostic criteria require validation in prospective clinical trials. AP-CML defined only by clonal cytogenetic evolution on imatinib therapy is associated with a better prognosis than AP-CML defined by clonal cytogenetic evolution and additional features of progression.^{43,72}

The International Bone Marrow Transplant Registry (IBMTR) criteria define blast phase as the presence of ≥30% myeloblasts in the blood, bone



NCCN Guidelines Version 2.2022

Chronic Myeloid Leukemia

marrow, or both, or as the presence of extramedullary disease ([CML-B](#)).⁷³ Any increase in lymphoblasts should be concerning for nascent lymphoid blast phase disease. IBMTR criteria were used in most of the clinical trials leading to the approval of TKIs, and is best aligned with prognostication systems derived from these studies. The WHO diagnostic criteria (presence of $\geq 20\%$ blast cells in the peripheral blood or bone marrow, the presence of extramedullary blast proliferation, and large foci or clusters of blasts in the bone marrow biopsy) were included in some reports.⁷¹ However, as clinical trials in the TKI era have almost uniformly utilized modified MD Anderson Cancer Center or IBMTR criteria, the use of WHO criteria for staging is disfavored.

Flow cytometry to determine cell lineage, mutational analysis, and human leukocyte antigen (HLA) testing, if considering allogeneic hematopoietic cell transplant (HCT), are recommended for patients with advanced phase CML.

Management of Chronic Phase CML

Primary Treatment ([CML-2](#))

Long-term efficacy data from randomized phase III studies for first-line TKI therapy in patients with newly diagnosed CP-CML are summarized in [Table 1](#).⁷⁴⁻⁷⁷ In summary, 1) all TKIs are highly effective in newly diagnosed CP-CML, with long-term OS approaching that of aged-matched controls; 2) second-generation TKIs, compared to imatinib, generally result in faster cytogenetic and molecular responses, with less progression to advanced phase CML; and 3) as of yet, in randomized clinical trials, there are no significant differences in OS in patients who initiate imatinib versus a second-generation TKI (dasatinib, nilotinib, and bosutinib).

The selection of first-line TKI therapy (bosutinib, dasatinib, imatinib, or nilotinib) in a given patient should be based on the risk score, toxicity

profile, patient's age, ability to tolerate therapy, and the presence of comorbid conditions. Allogeneic HCT is no longer recommended as a first-line treatment for patients with CP-CML.

Imatinib 800 mg is not recommended as initial therapy, given the recent data showing superior efficacy of second-generation TKIs in newly diagnosed CP-CML. Data from randomized phase III studies that have evaluated high-dose imatinib as first-line therapy for CP-CML suggest that imatinib 800 mg was not associated with lower rates of disease progression than imatinib 400 mg in any of these studies, despite improved early responses ([Table 2](#)).⁷⁸⁻⁸⁰ Imatinib 800 mg was also associated with higher rates of dose interruption, reduction, or discontinuation due to grade 3 or 4 adverse events in all of the studies. However, patients who were able to tolerate the higher dose of imatinib achieved higher response rates than those receiving standard-dose imatinib.⁸¹

Clinical Considerations for the Selection of First-Line Therapy

Risk Stratification

Imatinib (400 mg daily) and second-generation TKIs (bosutinib [400 mg daily], dasatinib [100 mg once daily], and nilotinib [300 mg twice daily]) are all appropriate options for first-line TKI therapy for patients with CP-CML across all risk scores.⁷⁴⁻⁷⁷

Disease progression is more frequent in patients with intermediate- or high-risk score, and prevention of disease progression to AP-CML or BP-CML is the primary goal of TKI therapy in patients with CP-CML. Second-generation TKIs are associated with lower risk of disease progression than imatinib and are therefore preferred for patients with an intermediate- or high-risk Sokal or Euro score.

Second-generation TKIs also result in quicker molecular responses and higher rates of major molecular response (MMR; $\leq 0.1\%$ *BCR-ABL1* IS)



NCCN Guidelines Version 2.2022

Chronic Myeloid Leukemia

and deep molecular response (DMR; MR4.5 [$\leq 0.0032\%$ *BCR-ABL1* IS]) in patients with CP-CML across all risk scores ([Table 3A](#)), which may facilitate subsequent discontinuation of TKI therapy in selected patients.⁷⁵⁻⁷⁷ In the ENESTnd study, nilotinib was also associated with lower rates of disease progression and higher PFS rates in patients with intermediate- and high-risk score ([Table 3B](#)).⁷⁶

Therefore, second-generation TKIs may be preferred over imatinib for younger patients, particularly women since the achievement of a deep and rapid molecular response may allow for eventual safe interruption of TKI therapy for fertility purposes. Imatinib may be preferred for older patients with comorbidities, especially cardiovascular comorbidities.

Toxicity Profile

All the TKIs are well tolerated. Since bosutinib, dasatinib, and nilotinib have very good efficacy in the upfront setting, differences in their potential toxicity profiles may inform the selection of a specific TKI as initial therapy. Nilotinib or bosutinib may be preferred for patients with a history of lung disease or deemed to be at risk of developing pleural effusions. Dasatinib or bosutinib may be preferred in patients with a history of arrhythmias, heart disease, pancreatitis, or hyperglycemia.

Adverse events of first-line TKI therapy in patients with CP-CML reported in phase III randomized studies are discussed below and are also summarized in [Table 4](#). See [CML-F](#) for the management of toxicities associated with TKI therapy.

Bosutinib

In the BFORE study, diarrhea, increased alanine aminotransferase (ALT), and aspartate aminotransferase (AST) were more common with bosutinib whereas muscle spasms and peripheral edema were more common with imatinib.⁷⁷ Grade 3/4 thrombocytopenia was higher with bosutinib and grade 3/4 neutropenia was higher with imatinib. Grade 3/4 anemia was

similar in both groups. Discontinuation of therapy due to drug-related adverse events occurred in 14% of patients in the bosutinib group compared to 11% in the imatinib group. Increased ALT (5%) and increased AST (2%) were the most common adverse events leading to discontinuation of bosutinib. However, there were no hepatotoxicity-related fatalities during the study.

Dasatinib

In the DASISION study, the incidences of grade 3/4 hematologic toxicities (anemia, neutropenia, and thrombocytopenia) were higher for dasatinib than imatinib.⁷⁵ Nonhematologic adverse events such as muscle spasms, peripheral edema, and hypophosphatemia were more frequent with imatinib. Discontinuation of therapy because of drug-related adverse events occurred in 16% and 7% of patients in the dasatinib and imatinib arms, respectively. Dasatinib is associated with significant but reversible inhibition of platelet aggregation that may contribute to bleeding in some patients, especially if accompanied by thrombocytopenia.⁸²

Pleural effusion was also more common with dasatinib (28% in the DASISION study compared to <1% with imatinib and 33% in a dose optimization study) and age has been identified as a significant risk factor for the development of pleural effusion.⁸³ The occurrence of pleural effusion is significantly reduced with dasatinib 100 mg once daily compared with 70 mg twice daily. Patients with prior cardiac history, with hypertension, and receiving dasatinib 70 mg twice daily are at increased risk of developing pleural effusions.⁸⁴ Close monitoring and timely intervention are necessary for patients at risk of developing pleural effusions.

Largely reversible pulmonary arterial hypertension has been reported as a rare but serious side effect of dasatinib.⁸⁵⁻⁸⁷ In the DASISION study, pulmonary hypertension was reported in 5% of patients treated with



NCCN Guidelines Version 2.2022

Chronic Myeloid Leukemia

dasatinib compared to <1% of patients treated with imatinib.⁷⁵ Evaluation for signs and symptoms of underlying cardiopulmonary disease prior to initiating and during treatment with dasatinib is recommended. If pulmonary arterial hypertension is confirmed, dasatinib must be permanently discontinued.

The recommended starting dose of dasatinib is 100 mg once daily for patients with newly diagnosed CP-CML. Long-term follow-up results of a single-arm study in a small cohort of patients suggest that dasatinib 50 mg once daily may have similar efficacy.⁸⁸ Treatment interruption of dasatinib at 100 mg once daily and reintroduction at a lower dose (40 mg twice daily or 60 mg once daily) has been shown to be effective for patients with intolerance to dasatinib at 100 mg once daily.^{89,90} Dasatinib at 50 mg (20 mg with careful monitoring in selected patients) should be considered for patients with clinically significant intolerance to dasatinib 100 mg once daily to avoid serious adverse events necessitating the discontinuation of dasatinib (eg, pleural effusion, myelosuppression). However, the minimum effective dasatinib dose has not been established in randomized clinical trials.

Imatinib

Chronic fatigue (often correlated with musculoskeletal pain and muscular cramps) is a major factor reducing quality of life.⁹¹ Hypophosphatemia and decrease in bone mineral density have been noted in a small group of patients, suggesting that monitoring bone health should be considered for patients taking imatinib.^{92,93} Skin hypopigmentation has also been reported as a side effect of imatinib and is reversible upon discontinuation or dose reduction.^{94,95} Reversible renal dysfunction with prolonged use of imatinib has also been reported.⁹⁶

Nilotinib

In the ENESTnd study, rates of nonhematologic adverse events such as nausea, diarrhea, vomiting, muscle spasm, and peripheral edema of any grade were higher for patients receiving imatinib. Conversely, rash and headache were more common with nilotinib.⁷⁶ Grade 3 or 4 neutropenia was more frequently observed in the imatinib group, whereas thrombocytopenia and anemia were similar in both groups. Electrolyte abnormalities and elevations in lipase, glucose, and bilirubin were more frequent with nilotinib than with imatinib. Patients with a previous history of pancreatitis may be at greater risk of elevated serum lipase. The overall incidences of adverse events leading to discontinuation of therapy were comparable in the nilotinib 300 mg twice-daily and imatinib arms (12% and 14%, respectively) and slightly higher in the nilotinib 400 mg twice-daily arm (20%).

Nilotinib labeling contains a black box warning regarding the risk of QT interval prolongation, and sudden cardiac death has been reported in patients receiving nilotinib.⁸⁷ QT interval prolongation could be managed with dose reduction. Electrolyte abnormalities should be corrected prior to initiation of treatment with nilotinib and electrolytes should be monitored periodically. Drugs that prolong QT interval should be avoided. Electrocardiogram (ECG) should be obtained to monitor the QT interval at baseline, 7 days after initiation of nilotinib, and periodically thereafter, as well as following any dose adjustments. Patients with cardiovascular risk factors should be referred to a cardiologist.

Nilotinib is associated with an increased risk of peripheral arterial occlusive disease (PAOD).⁹⁷⁻¹⁰¹ Patients should be evaluated for pre-existing PAOD and vascular risk factors prior to initiating and during treatment with nilotinib. If PAOD is confirmed, nilotinib should be permanently discontinued.



NCCN Guidelines Version 2.2022

Chronic Myeloid Leukemia

The recommended starting dose of nilotinib is 300 mg twice daily for patients with newly diagnosed CP-CML. Limited data from small cohorts of patients suggest that lower doses of nilotinib (<600 mg per day) may be associated with better safety and efficacy than nilotinib 300 mg twice daily.¹⁰² However, as with dasatinib, the minimum effective dose of nilotinib has not been established in randomized clinical trials.

Management of Hematologic Toxicities of TKI Therapy

Cytopenias (anemia, neutropenia, and thrombocytopenia) should be managed with transient interruptions of TKI therapy and dose modifications. Please see the package insert for full prescribing information, available at www.accessdata.fda.gov, for the recommended dose modifications of specific TKI therapy.

Assessment of reticulocyte count, ferritin, iron saturation, vitamin B12, and folate and correction of nutritional deficiencies if present is recommended for patients with grade 3–4 anemia. Red blood cell transfusions are indicated in symptomatic patients. Myeloid growth factor support can be used in combination with TKI therapy for the management of neutropenia.^{103,104} The use of erythropoiesis-stimulating agents (ESAs) did not impact survival or cytogenetic response rate, but was associated with a higher thrombosis rate in patients with CP-CML.¹⁰⁵ Recent guidelines from the U.S. Centers for Medicare & Medicaid Services (CMS) and the FDA do not support the use of ESAs in patients with myeloid malignancies.

Monitoring Response to TKI Therapy

Response to TKI therapy is determined by the measurement of hematologic (normalization of peripheral blood counts), cytogenetic (decrease in the number of Ph-positive metaphases using bone marrow cytogenetics), and molecular assessments (decrease in the amount of

BCR-ABL1 chimeric mRNA using qPCR). The criteria for hematologic, cytogenetic, and molecular response are summarized in [CML-D](#).

Conventional bone marrow cytogenetics is the standard method for monitoring cytogenetic responses, and many clinical trial response analyses were based on conventional bone marrow cytogenetics. With the advent of qPCR, bone marrow cytogenetic analyses to assess response are rarely performed at this time. If conventional bone marrow cytogenetics yield no analyzable metaphases, cytogenetic response can be evaluated by FISH, preferably with a dual color probe to minimize false-positive rates. FISH and cytogenetic results are correlated, but not superimposable.¹⁰⁶⁻¹⁰⁸ Although some investigators have reported that interphase FISH can be used to monitor complete cytogenetic response (CCyR), endpoints for TKI failure have not been defined on the basis of FISH analysis.^{109,110} The panel feels that FISH has been inadequately studied for monitoring response to TKI therapy and is not generally recommended for monitoring response if conventional cytogenetics or qPCR are available.

qPCR is the only tool capable of monitoring responses after the patient has achieved CCyR, since *BCR-ABL1* transcripts typically remain detectable after CCyR is achieved. A major advantage of qPCR is the strong correlation between the results obtained from the peripheral blood and the bone marrow, allowing for molecular monitoring without bone marrow aspirations.^{111,112}

Standardization of Molecular Monitoring Using the International Scale

In the IS, the standardized baseline (defined as the average expression of *BCR-ABL1* transcripts in 30 patients with untreated CML enrolled in the IRIS trial) is set to 100%. Molecular response is expressed as log-reduction from 100%. For example, a 2-log reduction or greater ($\leq 1\%$ *BCR-ABL1* IS; MR2.0) generally correlates with CCyR and a ≥ 3 -log reduction ($\leq 0.1\%$ *BCR-ABL1* IS) is referred to as MMR or MR3.0.^{14,113,114}

DMR is defined by the assay's level of sensitivity [$\leq 0.01\%$ *BCR-ABL1* (IS), MR4.0; $\leq 0.0032\%$ *BCR-ABL1* (IS), MR4.5].¹¹⁵ The sensitivity of a qPCR assay depends not only on the performance of the assay, but also on the quality of a given sample.

As such, the term “complete molecular response” to denote undetectable *BCR-ABL1* transcripts (a negative qPCR test) should be abandoned, as it may refer to very different levels of response, dependent on the quality of the sample. Laboratories can use their individual assays, but the *BCR-ABL1* transcripts obtained in a given laboratory should be converted to the IS by applying a laboratory-specific conversion factor (CF).^{14,116}

Recommendations for Monitoring Response to TKI Therapy

qPCR (IS) is the preferred method to monitor response to TKI therapy. qPCR assays with a sensitivity of ≥ 4.5 -log reduction from the standardized baseline are recommended for the measurement of *BCR-ABL1* transcripts (CML-C). In patients with prolonged myelosuppression who may not be in complete hematologic response (CHR) due to persistent cytopenias or unexplained drop in blood counts during therapy, bone marrow cytogenetics is indicated to confirm response to TKI therapy and exclude other pathology, such as MDS or the presence of chromosomal abnormalities other than Ph.

Monitoring with qPCR (IS) every 3 months is recommended for all patients after initiating TKI therapy, including those who meet response milestones at 3, 6, and 12 months ($\leq 10\%$ *BCR-ABL1* IS at 3 and 6 months, $\leq 1\%$ *BCR-ABL1* IS at 12 months, and $\leq 0.1\%$ *BCR-ABL1* IS at >12 months). After CCyR ($\leq 1\%$ *BCR-ABL1* IS) has been achieved, molecular monitoring is recommended every 3 months for 2 years and every 3 to 6 months thereafter.

Frequent molecular monitoring with qPCR (IS) can help to identify non-adherence to TKI therapy early in the treatment course.¹¹⁷ Since

adherence to TKI therapy is associated with better clinical outcomes, frequent molecular monitoring is essential if there are concerns about the patient's adherence to TKI therapy. In patients with deeper molecular responses (MMR and better) and who are compliant with TKI therapy, the frequency of molecular monitoring can be reduced, though the optimal frequency is unknown. Molecular monitoring of response to TKI therapy more frequently than every 3 months is not presently recommended.

Prognostic Significance of Cytogenetic and Molecular Response

Early molecular response (EMR; $\leq 10\%$ *BCR-ABL1* IS at 3 and 6 months) after first-line TKI therapy has emerged as an effective prognosticator of favorable long-term PFS and OS (Table 5).^{75,76,80,118} Some reports suggest that EMR at 3 months has a superior prognostic value and support the use of early intervention strategies based on the *BCR-ABL1* transcript level at 3 months.^{119,120} However, other studies yielded partially conflicting results regarding the predictive value of *BCR-ABL1* transcripts at 3 months.¹²¹ From a practical perspective, it is important to consider these data points within the clinical context. For instance, if *BCR-ABL1* transcript level is minimally above the 10% cutoff (11% at 3 months), it is reasonable to re-assess at 6 months before considering major changes to the treatment strategy.

Quite recently, studies have suggested that the rate of decline in *BCR-ABL1* transcripts correlates with longer-term response.¹²²⁻¹²⁴ Among patients with $>10\%$ *BCR-ABL1* IS after 3 months of treatment with imatinib, those with a faster decline in *BCR-ABL1* (*BCR-ABL1* halving time <76 days) had a superior outcome compared to those with a slower decline (4-year PFS rate was 92% vs. 63%, respectively).¹²² In the German CML IV study, lack of a half-log reduction of *BCR-ABL1* transcripts at 3 months was associated with a higher risk of disease progression on imatinib therapy.¹²³ The results of the D-First study also showed that in patients treated with dasatinib, *BCR-ABL1* halving time of



NCCN Guidelines Version 2.2022

Chronic Myeloid Leukemia

≤14 days was a significant predictor of MMR by 12 months and DMR (MR4.0; ≤0.01% *BCR-ABL1* IS) by 18 months.¹²⁴

Achievement of CCyR or ≤1% *BCR-ABL1* IS within 12 months after first-line TKI therapy is an established prognostic indicator of long-term survival.^{125,126} In the IRIS study, the estimated 6-year PFS rate was 97% for patients achieving a CCyR at 6 months compared to 80% for patients with no cytogenetic response at 6 months.¹²⁵ In an analysis of patients with newly diagnosed CP-CML treated with imatinib or second-generation TKIs, the 3-year event-free survival (EFS) and OS rates were 98% and 99% for patients who achieved CCyR at 12 months compared to 67% and 94% in patients who did not achieve a CCyR.¹²⁶

MMR (≤0.1% *BCR-ABL1* IS) as a predictor of PFS and OS has also been evaluated in several studies.^{111,127-133} In all of these studies, the analyses were done for different outcomes measures at multiple time points, but failed to adjust for multiple comparisons, thereby reducing the validity of the conclusions. The general conclusion from these studies is that the achievement of MMR is associated with durable long-term cytogenetic remission and lower rate of disease progression, but MMR is not a significant predictor of superior OS in patients who are in stable CCyR. Importantly, with longer follow-up, CCyR becomes an ever-stronger indicator of MMR, reducing the added prognostic value of MMR.

Although the CML IV study showed that MR4.5 (≤0.0032% *BCR-ABL1* IS) at 4 years was associated with a significantly higher OS (independent of therapy) than MR2.0 (≤1% *BCR-ABL1* IS which corresponds to CCyR), this study demonstrated no significant differences in OS in patients who achieved MMR (≤0.1% *BCR-ABL1* IS) and those who achieved MR2.0 (≤1% *BCR-ABL1* IS).¹³²

The absence of MMR in the presence of a CCyR is therefore not considered a treatment failure. While some investigators have reported that dose escalation of imatinib might benefit patients in CCyR with no

MMR,¹³⁴ there are no randomized studies to show that a change of therapy would improve survival, PFS, or EFS in this group of patients.¹³⁵ However, the achievement of MMR (≤0.1% *BCR-ABL1* IS) at 12 months is associated with a very low probability of subsequent loss of response and a high likelihood of achieving a subsequent DMR (MR4.0; ≤0.01% *BCR-ABL1* IS), which may facilitate discontinuation of TKI therapy.^{32,133} In view of the ongoing evolution of treatment goals (OS vs. treatment-free remission [TFR]), expert panels have emphasized the importance of joint decision-making between patient and provider, particularly in ambiguous situations.¹³⁶

Response Milestones After First-Line TKI Therapy (CML-3)

The most important goals of TKI therapy are to prevent disease progression to AP-CML or BP-CML and to achieve either MR2.0 (≤1% *BCR-ABL1* IS, which corresponds to CCyR) or MMR (≤0.1% *BCR-ABL1* IS) within 12 months after first-line TKI therapy. The guidelines emphasize that achievement of response milestones must be interpreted within the clinical context, before making drastic changes to the treatment strategy, especially in ambiguous situations.

The panel has included ≤10% *BCR-ABL1* IS at 3 and 6 months after initiation of first-line TKI therapy as a response milestone, since the achievement of EMR after first-line TKI therapy is an effective prognosticator of favorable long-term PFS. Achievement of >0.1%–1% *BCR-ABL1* IS (≤1% *BCR-ABL1* IS, which correlates with CCyR) is considered the optimal response milestone at 12 months if the goal of therapy in an individual patient is long-term survival, whereas the achievement of MMR (≤0.1% *BCR-ABL1* IS) at 12 months should be considered as the optimal response milestone if the treatment goal in an individual patient is TFR. Patients who achieve these response milestones are considered to have TKI-sensitive disease, and continuation of the



NCCN Guidelines Version 2.2022

Chronic Myeloid Leukemia

same dose of TKI and assessment of *BCR-ABL1* transcripts with qPCR (IS) every 3 months is recommended for this group of patients.

In patients with a >10% *BCR-ABL1* IS at 3 months and >1% *BCR-ABL1* IS at 12 months, clinical judgment should be used, considering problems with adherence (which can be common given drug toxicity at initiation of therapy), rate of decline in *BCR-ABL1* (the faster, the better), and how far from the cutoff the *BCR-ABL1* value falls. That being said, failure to achieve ≤10% *BCR-ABL1* IS at 3 months or ≤1% *BCR-ABL1* IS at 12 months is associated with a higher risk for disease progression. Patients with >10% *BCR-ABL1* at 3 months or >1% *BCR-ABL1* at 12 months can continue the same dose of dasatinib, nilotinib, or bosutinib for another 3 months. *BCR-ABL1* mutational analysis and evaluation for allogeneic HCT should be considered. Bone marrow cytogenetics should be considered to assess for major cytogenetic response (MCyR) at 3 months or CCyR at 12 months.

In patients with >0.1%–1% *BCR-ABL1* IS at 12 months, shared decision-making is recommended depending on the goal of therapy in individual patients (longer-term survival vs. TFR). As discussed before, MMR at 12 months is associated with lower rate of disease progression and a higher likelihood of achieving DMR, which is a prerequisite for TFR. Switching to a second-generation TKI from imatinib might be considered to increase the probability of achieving MMR (≤0.1% *BCR-ABL1* IS) at 12 months. However, it is also associated with increased toxicity. Referral to specialized CML centers and/or enrollment in a clinical trial should be considered.

Patients with >10% *BCR-ABL1* IS at 6 and 12 months are considered to have TKI-resistant disease. Evaluation for allogeneic HCT (that is, a discussion with a transplant specialist, which might include HLA testing) is recommended. Alternate treatment options should be considered as described below.

Second-line Therapy

Long-term efficacy data from phase II/III studies on second-line and subsequent TKI therapy for CP-CML are summarized in [Table 6](#).¹³⁷⁻¹⁴⁰

EMR (≤10% *BCR-ABL1* IS at 3 and 6 months) after second-line TKI therapy with dasatinib or nilotinib has also been reported to be a prognosticator of OS and PFS ([Table 7](#)). Patients who do not achieve cytogenetic or molecular responses at 3, 6, or 12 months after second-line and subsequent TKI therapy should be considered for alternative therapies or allogeneic HCT if deemed eligible.

Management of Patients with Inadequate Response to Imatinib

Switching to an alternate TKI is recommended for patients with disease that is resistant to imatinib 400 mg daily. Dasatinib, nilotinib, and bosutinib, which are more potent than imatinib *in vitro* and retain activity against many of the imatinib-resistant *BCR-ABL1* kinase domain mutants except T315I, are effective treatment options for patients with CP-CML intolerant or resistant to imatinib.¹³⁷⁻¹³⁹

Dose escalation of imatinib up to 800 mg daily has been shown to overcome some cases of primary resistance and is particularly effective for cytogenetic relapse in patients who had achieved cytogenetic response with imatinib 400 mg daily, although the duration of responses has typically been short.¹⁴¹⁻¹⁴⁴ However, it is unlikely to benefit patients with hematologic failure or those who never had a cytogenetic response with imatinib 400 mg daily. In patients with inadequate response to imatinib 400 mg, switching to nilotinib has been shown to result in higher rates of cytogenetic and molecular response than dose escalation of imatinib.^{145,146} In the TIDEL-II study, the cohort of patients with >10% *BCR-ABL1* IS at 3 months after imatinib 400 mg who were switched directly to nilotinib had higher rates of MMR and complete molecular response (CMR) at 12 months (but not at 24 months) than the cohort of patients who received



NCCN Guidelines Version 2.2022

Chronic Myeloid Leukemia

dose escalation of imatinib before switching to nilotinib.¹⁴⁵ Although dose escalation of imatinib has been shown to be beneficial for patients in CCyR with no MMR, no randomized studies have shown that a change of therapy would improve PFS or EFS in this group of patients.^{134,135}

Management of Patients with Inadequate Response to Dasatinib, Nilotinib, or Bosutinib

Bosutinib has demonstrated activity in patients with CP-CML resistant/intolerant to multiple TKIs (imatinib, dasatinib, and nilotinib; [Table 6](#)).^{147,148} Ponatinib was initially approved as a treatment option for patients with a T315I mutation and/or for patients for whom no other TKI is indicated based on the results of the PACE trial ([Table 6](#)).¹⁴⁰

In the PACE trial, serious arterial occlusive events (cardiovascular, cerebrovascular, and peripheral vascular) and venous thromboembolic events occurred in 31% and 6% of patients, respectively.¹⁴⁰ Cardiovascular, cerebrovascular, and peripheral arterial occlusive events were reported in 16%, 13%, and 14% of patients, respectively. In an analysis of cardiovascular, arterial, and thrombotic adverse events associated with front-line TKI therapy in prospective clinical trials, the incidence of cardiovascular adverse events was highest among patients treated with ponatinib and those with preexisting cardiovascular risk factors.¹⁴⁹ The increased incidences of arterial occlusive events among patients treated with ponatinib were also confirmed in another multicenter real-life study.¹⁵⁰

The recommended initial dose of ponatinib was 45 mg once daily, the maximum tolerated dose determined in a phase 1 dose-escalation study.¹⁵¹ As high-dose intensity of ponatinib is associated with increased risk of adverse events, dose modifications may be necessary to prevent or manage adverse events.¹⁵² Recent reports suggest that substantial responses can be observed at lower dose levels (30 mg or 15 mg) with

decreased incidence of cardiovascular events; the rates at which MCyR and MMR were maintained were independent of the dose reductions.^{140,153}

The OPTIC trial (results of which have only been published as an abstract) evaluated the safety and efficacy of response-adjusted dosing regimen of ponatinib in patients with CP-CML (283 patients with CP-CML resistant or intolerant to ≥ 2 TKIs or with a T315I mutation were randomized to receive 3 different doses of ponatinib (once daily): 45 mg (cohort A), 30 mg (B), and 15 mg (C). Doses were reduced to 15 mg after achievement of *BCR-ABL1* (IS) $\leq 1\%$. After a median follow-up of 12 months, *BCR-ABL1* (IS) $\leq 1\%$ was achieved in 39%, 27% and 26% of patients in the 3 different dose cohorts respectively.

Based on the results of OPTIC trial, the FDA has approved a response-based dosing regimen for ponatinib for patients with CP-CML with resistance or intolerance to at least two prior kinase inhibitors. The recommended initial dose is 45 mg once daily with a reduction to 15 mg upon achievement of *BCR-ABL1* (IS) $\leq 1\%$. The ongoing OPTIC trial is also evaluating the safety and efficacy of lower initial doses of ponatinib (30 mg and 15 mg).

The ponatinib labeling contains a black box warning regarding arterial occlusive events, venous thromboembolic events, heart failure, and hepatotoxicity. Cardiovascular risk factors (eg, diabetes mellitus, hypertension, hyperlipidemia, smoking, estrogen use) should be identified and controlled before starting ponatinib. Patients should be monitored for high blood pressure, evidence of arterial occlusive or thromboembolic events, and reduced cardiac function.¹⁵⁴ Ponatinib should be interrupted or stopped immediately for vascular occlusion and for new or worsening heart failure. Patients with cardiovascular risk factors should be referred to a cardiologist.



NCCN Guidelines Version 2.2022

Chronic Myeloid Leukemia

Switching to an alternate TKI (other than imatinib) could be considered for patients with disease that is resistant to dasatinib, nilotinib, or bosutinib. However, there is no clear evidence to support that switching to alternate TKI therapy would improve long-term clinical outcome for this group of patients. The use of an alternate second-generation TKI after treatment failure with *two* prior TKIs, including a second-generation TKI, is not associated with durable responses except in occasional patients with CP-CML.¹⁵⁵

Omacetaxine is a treatment option for patients with CP-CML resistant or intolerant to ≥ 2 TKIs including those with a T315I mutation.^{156,157} In the CML 202 study, among 62 evaluable patients with CP-CML resistant to prior TKI therapy and T315I mutation, after a median follow-up of 19 months, MCyR, CCyR, and MMR rates were 23%, 16%, and 17%, respectively, and the T315I clone declined to below detection limits in 61% of patients.¹⁵⁶ The median PFS was 8 months and the median OS had not yet been reached. In the cohort of 46 patients with CP-CML resistant or intolerant to ≥ 2 TKIs (CML 203 study), after a median follow-up of 19 months, the MCyR and CCyR rates were 22% and 4%, respectively. The median PFS and OS were 7 months and 30 months, respectively.¹⁵⁷ The response rates and survival outcomes, however, were substantially lower than that observed with ponatinib in the PACE trial in this patient population ([Table 6](#); the estimated 5-year PFS rate was 52% for patients with CP-CML resistant or intolerant to ≥ 2 TKIs and 50% for those with a T315I mutation).¹⁴⁰ Omacetaxine had an acceptable toxicity profile and the most common grade 3/4 adverse events were thrombocytopenia (67%), neutropenia (47%), and anemia (37%).

Clinical Considerations for the Selection of Second-Line Therapy

BCR-ABL1 kinase domain mutation analysis (see below), evaluation of drug interactions, and compliance to therapy are recommended prior to the initiation of second-line TKI therapy.

Drug Interactions

Bosutinib, dasatinib, imatinib, nilotinib, and ponatinib are metabolized in the liver by cytochrome P450 (CYP) enzymes, and concomitant use of drugs that induce or inhibit CYP3A4 or CYP3A5 enzymes may alter the therapeutic effect of TKIs.^{158,159} Drugs that are CYP3A4 or CYP3A5 inducers may decrease the therapeutic plasma concentration of TKIs, whereas CYP3A4 inhibitors and drugs that are metabolized by the CYP3A4 or CYP3A5 enzyme might result in increased plasma levels of TKIs. In addition, imatinib is also a weak inhibitor of the CYP2D6 and CYP2C9 isoenzymes and nilotinib is a competitive inhibitor of CYP2C8, CYP2C9, CYP2D6, and UGT1A1, potentially increasing the plasma concentrations of drugs eliminated by these enzymes.

Drug interactions between TKIs and some of the most commonly used drugs and supplements are summarized in [CML-F](#). Concomitant use of drugs that are metabolized by these enzymes requires caution, and appropriate alternatives should be explored to optimize treatment outcome. If coadministration cannot be avoided, dose modification should be considered.

Adherence to Therapy

Treatment interruptions and non-adherence to therapy may lead to undesirable clinical outcomes.¹⁶⁰⁻¹⁶² In the ADAGIO study, non-adherence to imatinib was associated with poorer response. Patients with suboptimal response missed significantly more imatinib doses (23%) than did those with optimal response (7%).¹⁶⁰ Adherence to imatinib therapy has been identified as the only independent predictor for achieving CMR on standard-dose imatinib.¹⁶¹ Poor adherence to imatinib therapy has also been identified as the most important factor contributing to cytogenetic relapse and imatinib failure.¹⁶² Patients with adherence of $\leq 85\%$ had a higher probability of losing CCyR at 2 years than those with adherence of $>85\%$ (27% and 2%, respectively). Poor adherence to therapy has also



NCCN Guidelines Version 2.2022

Chronic Myeloid Leukemia

been reported in patients receiving dasatinib and nilotinib following imatinib failure.^{163,164}

Patient education on adherence to therapy and close monitoring of patient's adherence is critical to achieving optimal responses. In a significant proportion of patients with TKI-induced toxicities, responses have been observed with doses well below their determined maximum tolerated doses.¹⁶⁵ Short interruptions or dose reductions, when medically necessary, may not have a negative impact on disease control or other outcomes. Adequate and appropriate management of side effects and scheduling appropriate follow-up visits to review side effects may be helpful to improve patient adherence to therapy.¹⁶⁶ Switching to an alternate TKI because of intolerance might be beneficial for selected patients with acute grade 3/4 non-hematologic toxicities or in those with chronic, low-grade nonhematologic toxicities that are not manageable with adequate supportive care measures.¹⁶⁷⁻¹⁶⁹

Resistance to TKI Therapy

Aberrant expressions of drug transporters¹⁷⁰⁻¹⁷² and plasma protein binding of TKI¹⁷³⁻¹⁷⁵ could contribute to primary resistance by altering the intracellular and plasma concentration of TKI.

Pretreatment levels of organic cation transporter 1 (OCT1) have been reported as the most powerful predictor of response to imatinib.¹⁷⁶ On the other hand, cellular uptake of dasatinib or nilotinib seems to be independent of OCT1 expression, suggesting that patients with low OCT1 expression might have better outcomes with dasatinib or nilotinib than with imatinib.¹⁷⁷⁻¹⁸⁰

Monitoring imatinib plasma levels may be useful in determining patient adherence to therapy. However, there are no data to support that change of therapy based on plasma imatinib levels will affect treatment

outcomes, and assays that measure plasma levels of imatinib are not widely available.

BCR-ABL1 Kinase Domain Mutation Analysis

Point mutations in the BCR-ABL1 kinase domain are a frequent mechanism of secondary resistance to TKI therapy and are associated with poor prognosis and higher risk of disease progression.¹⁸¹⁻¹⁸⁶ Among the BCR-ABL1 kinase domain mutations, T315I confers complete resistance to imatinib, dasatinib, nilotinib, and bosutinib.^{187,188} The T315A, F317L/I/V/C, and V299L mutants are resistant to dasatinib and E255K/V, F359V/C, and Y253H mutants are resistant to nilotinib.¹⁸⁹⁻¹⁹² E255K/V, F359C/V, Y253H, and T315I mutants are most commonly associated with disease progression and relapse.^{192,193} Bosutinib has demonstrated activity in patients with BCR-ABL1 mutants resistant to dasatinib (F317L) and nilotinib (Y253H, E255K/V, and F359C/I/V).¹⁴⁷ However, bosutinib has minimal activity against F317L mutant while *in vitro* studies suggest that F317L is highly sensitive to nilotinib.^{190,192,194} Nilotinib may be preferred over bosutinib in patients with F317L mutation. T315I, G250E, and V299L mutants are resistant to bosutinib.¹⁴⁷ Ponatinib is active against *BCR-ABL1* mutants resistant to dasatinib or nilotinib, including E255V, Y253H, and F359V, in addition to T315I.¹⁴⁰ Response rates to TKI therapy based on BCR-ABL mutation status are listed in [Table 8](#).

BCR-ABL1 compound mutations (variants containing ≥2 mutations within the same *BCR-ABL1* allele that presumably arise sequentially) confer different levels of resistance to TKI therapy, and T315I-inclusive compound mutants confer the highest level of resistance to all TKIs, including ponatinib.^{195,196} In a more recent study that used NGS to detect low-level and *BCR-ABL1* compound mutations in 267 patients with heavily pretreated CP-CML from the PACE trial, no compound mutation was identified that consistently conferred resistance to ponatinib, suggesting



NCCN Guidelines Version 2.2022

Chronic Myeloid Leukemia

that such compound mutations are uncommon following treatment with bosutinib, dasatinib, or nilotinib for CP-CML.¹⁹⁷

BCR-ABL1 kinase domain mutational analysis is helpful in the selection of subsequent TKI therapy for patients with inadequate initial response to first-line or second-line TKI therapy.¹⁹⁸ The guidelines recommend *BCR-ABL1* mutational analysis for patients who do not achieve response milestones, for those with any sign of loss of response (hematologic or cytogenetic relapse), and if there is a 1-log increase in *BCR-ABL1* level with loss of MMR. Treatment options based on BCR-ABL1 kinase domain mutation status are outlined on [CML-5](#).

BCR-ABL1 mutational analysis provides additional guidance in the selection of subsequent TKI therapy only in patients with identifiable mutations. In patients with no identifiable mutations, the selection of subsequent TKI therapy should be based on the toxicity profile of TKI, patient's age, ability to tolerate therapy, and the presence of comorbid conditions. Adverse events of second-line TKI therapy in patients with CP-CML are summarized in [Table 9](#).

BCR-ABL1-independent Mutations

Mutations in a variety of cancer-associated genes other than *BCR-ABL1* (eg, *ASXL1*, *RUNX1*, *IKZF1*, *TET1/2*, *IDH1/2*, *JAK2*, *DNMT3A/3B*, *EZH2*, *WT1*, *NPM1*, *NRAS*, *KRAS*, *CBL*, *BCOR*, *CREBBP*, and *TP53*) have been described in patients with CML at diagnosis and in patients with AP-CML or BP-CML.¹⁹⁹⁻²⁰⁶ *IKZF1* exon deletions and mutations in *ASXL1*, *RUNX1*, and *BCOR* genes were the most frequently described in advanced phase CML, while *IDH1/2* mutations were detected at a markedly lower frequency.^{205,206} *IKZF1* and *RUNX1* alterations, both involved in cell differentiation, were identified as important markers of disease progression from CP-CML to BP-CML.^{199,204} In one study that analyzed the mutation landscape of patients with CML using a panel of 92 genes associated with myeloid malignancy, the presence of mutations in genes involved in

epigenetic regulation pathways at diagnosis (eg, *ASXL1*, *BCOR*, *TET1/2*, *IDH1/2*, *DNMT3A/3B*, and *EZH2*) was associated with poor response to TKI therapy (CCyR at 12 months, $P = .02$; MMR at 24 months, $P = .04$; and MR4.5 at 36 months, $P = .03$) independent of other clinical factors.²⁰²

However, many of these studies did not indicate whether the patients had CP-CML at diagnosis, and the impact of mutations is also variable depending on whether they occur in Ph-positive or Ph-negative clones.²⁰⁵ Therefore, these results are not indicative of the frequency of mutations in cancer-associated genes in patients with CP-CML at diagnosis and results are not definitive.

Rising BCR-ABL1 Transcripts

Rising *BCR-ABL1* transcripts are associated with an increased likelihood of detecting BCR-ABL1 kinase domain mutations and cytogenetic relapse.²⁰⁷⁻²¹¹ In patients who had achieved very low levels of *BCR-ABL1* transcripts, emergence of BCR-ABL1 kinase domain mutations was more frequent in those who had >2-fold increase in *BCR-ABL1* transcripts compared to those with stable or decreasing *BCR-ABL1* transcripts.²⁰⁷ A serial rise has been reported to be more reliable than a single ≥ 2 -fold increase in *BCR-ABL1* transcripts.^{208,209} Among patients in CCyR with a ≥ 0.5 -log increase in *BCR-ABL1* transcripts on at least two occasions, the highest risk of disease progression was associated with loss of MMR and >1-log increase in *BCR-ABL1* transcripts.²⁰⁹

Rising transcript levels should prompt investigation of treatment adherence and reassessment of coadministered medications. The precise increase in *BCR-ABL1* transcripts that warrants a mutation analysis depends on the performance characteristics of the qPCR assay.²¹¹ Some labs have advocated a 2- to 3-fold range,^{130,210,211} while others have taken a more conservative approach (5 – 10-fold).²⁰⁹ Obviously, some common sense must prevail, since the amount of change in absolute terms depends on the level of molecular response. For example, a finding of any



NCCN Guidelines Version 2.2022

Chronic Myeloid Leukemia

BCR-ABL1 after achieving a DMR (MR4.5; $\leq 0.0032\%$ *BCR-ABL1* IS) is an infinite increase in *BCR-ABL1* transcripts. However, a change in *BCR-ABL1* transcripts from a barely detectable level to MR4.5 is clearly different from a 5-fold increase in *BCR-ABL1* transcripts after achieving MMR.

Currently there are no specific guidelines for changing therapy only based on rising *BCR-ABL1* levels as detected by qPCR, and it should be done only in the context of a clinical trial.

Discontinuation of TKI Therapy (CML-E)

The feasibility of discontinuation of TKI therapy (with close monitoring) in carefully selected patients who have achieved and maintained DMR ($\geq$ MR4.0; $\leq 0.01\%$ *BCR-ABL1* IS) for ≥ 2 or more years has been evaluated in several clinical studies. Limited longer-term follow-up data from the TKI discontinuation trials are summarized in [Table 10](#).

The possibility of TFR after discontinuation of imatinib was first evaluated in the Stop Imatinib (STIM1) study in 100 patients with undetectable *BCR-ABL1* transcripts for at least 2 years (5-log reduction in *BCR-ABL1* transcripts and undetectable minimal residual disease on qPCR with a sensitivity of ≥ 4.5 -log reduction from the standardized baseline).^{212,213} With a median follow-up of 77 months after discontinuation of imatinib, the molecular recurrence-free survival was 43% at 6 months and 38% at 60 months.²¹³ Other subsequent studies that have evaluated the discontinuation of imatinib have also reported similar findings.²¹⁴⁻²¹⁸

More recent studies have also confirmed the feasibility of TFR after discontinuation of dasatinib or nilotinib in patients with CP-CML who have achieved and maintained MR4.5 for 12 months after ≥ 2 years of TKI therapy in the first-line or second-line setting (TFR rates ranging from 44% to 54%; [Table 10](#)).²¹⁹⁻²²⁵ The feasibility of TFR following discontinuation of bosutinib or ponatinib has not yet been evaluated in

clinical studies. In the EURO-SKI study that evaluated TFR after discontinuation of any first-line TKI therapy (imatinib, dasatinib, or nilotinib) in eligible patients, the type of first-line TKI therapy did not significantly affect molecular relapse-free survival.²²³ Therefore, it is reasonable to assume that the likelihood of TFR following discontinuation would be similar irrespective of TKI in patients who have achieved and maintained DMR (MR4.0; $\leq 0.01\%$ *BCR-ABL1* IS) for ≥ 2 years.

The results of the RE-STIM study demonstrated the safety of a second TKI discontinuation after a first unsuccessful attempt.²²⁶ The rate of molecular relapse after the first TKI discontinuation attempt was the only factor significantly associated with outcome. The TFR rate at 24 months after second TKI discontinuation was higher for patients who remained in DMR within the first 3 months after the first TKI discontinuation (72% vs. 32% for other patients).

Approximately 40% to 60% of patients who discontinue TKI therapy after achieving DMR experience recurrence within 12 months of treatment cessation, in some cases as early as one month after discontinuation of TKI therapy. Resumption of TKI therapy immediately after recurrence results in the achievement of DMR in almost all patients.²¹²⁻²²⁴ TKI withdrawal syndrome (aggravation or new development of musculoskeletal pain and/or pruritus after discontinuation of TKI therapy) has been reported during the TFR period in some TKI discontinuation studies,^{217,220,221,224} and the occurrence of imatinib withdrawal syndrome was associated with a lower rate of molecular relapse in the KID study.²¹⁷

In the STIM study, molecular relapse (trigger to resume TKI therapy) was defined as positivity for *BCR-ABL1* transcripts by qPCR confirmed by a 1-log increase in *BCR-ABL1* transcripts between two successive assessments or loss of MMR at one point.^{212,213} The results of the A-STIM study showed that loss of MMR ($\leq 0.1\%$ *BCR-ABL1* IS) could be used as a practical criterion for restarting therapy. The estimated probability of MMR



NCCN Guidelines Version 2.2022

Chronic Myeloid Leukemia

loss was 35% at 12 months and 36% at 24 months after discontinuation of imatinib.²¹⁵ Several factors may help predict the risk of recurrence after discontinuation of TKI therapy (eg, a higher Sokal risk score, female gender, lower natural killer cell counts, suboptimal response or resistance to imatinib, duration of TKI therapy, and DMR prior to TKI discontinuation).^{212,213,217,219-224,227} However, only the duration of TKI therapy and DMR prior to TKI discontinuation therapy have been associated with TFR with a high level of consistency.^{212,217,223,224} In the EURO-SKI study, duration of treatment with imatinib (≥ 6 years) and duration of DMR (MR4.0 for 3 years) were significantly associated with MMR maintenance at 6 months after discontinuation of imatinib.²²³

Based on the available evidence from clinical studies that have evaluated the feasibility of TFR, the panel members feel that discontinuation of TKI therapy (with *close monitoring*) is feasible in carefully selected, consenting patients (in early CP-CML) who have achieved and maintained a DMR ($\geq$ MR4.0) for ≥ 2 years. Clinical studies that have evaluated the safety and efficacy of discontinuation of TKI have employed strict eligibility criteria and have mandated more frequent molecular monitoring than typically recommended for patients on TKI therapy. Access to a reliable qPCR (IS) with a sensitivity of detection of at least MR4.5 ($BCR-ABL1 \leq 0.0032\%$ IS) and the availability of test results within 2 weeks is one of the key requirements to monitor patients after discontinuation of TKI therapy and ascertain their safety.

The criteria for the selection of patients suitable for discontinuation of TKI therapy are outlined on [CML-E](#). The guidelines emphasize that discontinuation of TKI therapy outside of a clinical trial should be considered only if ALL the criteria included on the list are met. The panel acknowledges that more frequent molecular monitoring is essential following discontinuation of TKI therapy for the early identification of loss of MMR. Frequency of molecular monitoring has varied substantially

among different studies, and the optimal frequency of molecular monitoring in patients with a loss of MMR after discontinuation of TKI therapy has not been established. The panel recommendations for molecular monitoring in TFR phase are outlined on [CML-E](#).

Management of Advanced Phase CML

Imatinib has induced favorable hematologic and cytogenetic response rates in patients with AP-CML or BP-CML.²²⁸⁻²³² Dasatinib,²³³⁻²³⁵ nilotinib,^{236,237} bosutinib,²³⁸ and ponatinib¹⁴⁰ have demonstrated activity in imatinib-resistant or imatinib-intolerant AP-CML and/or BP-CML. Ponatinib is a treatment option for patients with T315I mutation or for patients for patients whom no other TKI is indicated.

Long-term follow-up data from phase II/III studies evaluating TKI therapy for disease progression to AP-CML and BP-CML are summarized in [Table 11](#) and [Table 12](#), respectively.

The efficacy of imatinib in combination with decitabine or cytarabine-based chemotherapy in AP-CML and myeloid BP-CML has been demonstrated in several small studies.²³⁹⁻²⁴² Hyper-CVAD in combination with imatinib or dasatinib is also effective for patients with lymphoid BP-CML, particularly when followed by allogeneic HCT.^{243,244}

A significant proportion of patients with AP-CML or BP-CML treated with TKI therapy achieve a MCyR but not a concomitant CHR because of persistent cytopenias, which in turn is associated with an inferior outcome.²⁴⁵ Omacetaxine has shown efficacy in patients with AP-CML that is resistant to multiple TKIs as well as for patients with a T315I mutation.²⁴⁶ Among the 51 patients with AP-CML, after a median follow-up of 16 months, major hematologic response (MaHR), CHR, and minor cytogenetic response were achieved or maintained in 37%, 29%, and 11% of patients, respectively.²⁴⁶ The MaHR rates were 55% and 58%, respectively, for patients with a history of a T315I mutation and for those



NCCN Guidelines Version 2.2022

Chronic Myeloid Leukemia

with confirmed T315I mutation at baseline. The median PFS and OS were 5 months and 18 months, respectively. As with CP-CML, the response rates and survival outcomes were lower than that observed with ponatinib in the PACE trial for patients with AP-CML ([Table 11](#); the estimated 5-year PFS rates were 19% for patients with CP-CML resistant or intolerant to ≥ 2 TKIs and 29% for those with T315I mutation. The corresponding 5-year OS rates were 48% and 52%, respectively).¹⁴⁰ The most common grade 3/4 hematologic adverse events were thrombocytopenia (51%), anemia (39%), neutropenia (20%), and febrile neutropenia (14%).

Treatment Considerations (CML-4)

Disease progression to AP-CML or BP-CML while on TKI therapy has a worse prognosis than *de novo* AP-CML or BP-CML. Participation in clinical trials and evaluation for allogeneic HCT is recommended for all patients with AP-CML and BP-CML. In patients with disease progression to AP-CML or BP-CML, the selection of TKI therapy is based on prior therapy and/or BCR-ABL1 kinase domain mutational analysis.

De novo AP-CML can often be initially managed like CP-CML with single-agent TKI followed by evaluation for allogeneic HCT.^{247,248} However, patients with disease progression from CP-CML to AP-CML while on a TKI therapy have a high rate of progression to BP-CML, with predictably poor survival. These patients should be considered for a clinical trial and/or allogeneic HCT. Treatment with a course of alternate TKI (not received before) can be beneficial as a “bridge” to allogeneic HCT in patients with disease progression to AP-CML. There is a lack of evidence for the definition of optimal response milestones on TKI therapy. Evaluation for allogeneic HCT should be considered if response milestones (recommended for CP-CML) are not achieved at 3, 6, and 12 months. Omacetaxine is also an option for patients with disease progression to AP-CML on TKI therapy.²⁴⁶

TKI in combination with chemotherapy (ALL-type chemotherapy for lymphoid BP-CML and AML-type chemotherapy for myeloid BP-CML) or steroids followed by allogeneic HCT is recommended for *de novo* BP-CML and disease progression to BP-CML.²⁴⁹

Central nervous system (CNS) involvement has been described in case reports of BP-CML.²⁵⁰⁻²⁵³ Lumbar puncture and CNS prophylaxis is recommended for lymphoid BP-CML. Documented CNS involvement in patients with lymphoid BP-CML should be managed according to the standard of care for AML or ALL. Dasatinib has been reported to cross the blood brain barrier and may represent the best TKI option for patients with CNS disease.²⁵⁴ TKI therapy has not been optimized for patients with CNS involvement.

Allogeneic Hematopoietic Cell Transplant

Allogeneic HCT is a potentially curative treatment for patients with CML. Ongoing advances in alternative donor sources (such as unrelated donors and cord blood), more accurate HLA testing for a stringent selection of unrelated matched donors, and the use of reduced-intensity conditioning regimens have improved outcomes following allogeneic HCT.²⁵⁵⁻²⁶¹

Allogeneic HCT is an appropriate treatment option for the very rare patients presenting with BP-CML at diagnosis, patients with disease that is resistant to TKIs, patients with progression to AP-CML or BP-CML while on TKI therapy, and patients with CML that is resistant and/or intolerant to all TKIs.²⁶²⁻²⁶⁵ Several studies have confirmed that prior TKI therapy does not compromise the outcome following allogeneic HCT or increase transplant-related toxicity.²⁶⁶⁻²⁷²

Disease phase, HLA matching, age and sex of the donor and recipient, and time from diagnosis to transplant have been identified as pretransplant risk factors.²⁷³ A low HCT comorbidity index is a prognostic indicator of lower non-relapse mortality and improved survival.²⁷⁴ The



NCCN Guidelines Version 2.2022

Chronic Myeloid Leukemia

disease phase at the time of transplant remains an important prognostic factor, and the survival outcomes following transplant are clearly better for patients in second chronic CP-CML compared to patients with AP-CML or BP-CML.²⁷⁵⁻²⁸⁰ Therefore, the potential use of allogeneic HCT must be tied to faithful monitoring of disease, since the major potential pitfall in delaying transplantation is “missing” the chronic phase interval.

Monitoring Response After Allogeneic HCT (CML-6)

BCR-ABL1 transcripts may persist for many years in patients after allogeneic HCT. The prognostic significance of *BCR-ABL1* positivity is influenced by the time of testing after allogeneic HCT. A positive qPCR assay for *BCR-ABL1* at ≥18 months after allogeneic HCT is associated with a lower risk of relapse than a positive qPCR assay for *BCR-ABL1* at 6 to 12 months after allogeneic HCT.²⁸¹⁻²⁸⁸ Early detection of *BCR-ABL1* transcripts after allogeneic HCT may be useful to identify patients who may be in need of alternative therapies before overt relapse occurs.

Management of Post-transplant Relapse (CML-6)

Donor lymphocyte infusion (DLI) is effective in inducing durable molecular remissions in the majority of patients with relapsed CML following allogeneic HCT, though it is more effective in patients with chronic phase relapse than advanced phase relapse.²⁸⁹⁻²⁹⁵ However, DLI is associated with complications such as graft-versus-host disease (GVHD), susceptibility to infections, and immunosuppression.²⁸⁹ Improvements in the methods of detecting *BCR-ABL1* transcripts to predict relapse, the development of reduced-intensity conditioning regimens, modified delivery of lymphocytes with the depletion of CD8+ cells, and the use of escalating cell dosage regimens have reduced the incidence of GVHD associated with DLI.²⁹⁶⁻³⁰⁰

Imatinib induces durable cytogenetic and molecular responses in the majority of patients relapsing with chronic and advanced phase CML

following allogeneic HCT, and the response rates are higher in patients with chronic phase relapse than advanced phase relapse.³⁰¹⁻³⁰⁸ Very limited data are available on the use of dasatinib and nilotinib in patients with post-transplant relapse.³⁰⁹⁻³¹² There are also data suggesting that the use of DLI in combination with imatinib may be more effective at inducing rapid molecular remissions than either modality alone.³¹³ Recent retrospective studies have shown that TKIs are superior to DLI alone or in combination with TKI for post-transplant relapse.^{314,315} However, these observations are yet to be confirmed in randomized trials. Post-transplant TKI therapy is also effective to prevent relapse following allogeneic HCT in high-risk patients.³¹⁶⁻³¹⁸

Patients who are in CCyR (qPCR-negative) should undergo regular qPCR monitoring (every 3 months for 2 years, then every 3–6 months thereafter). Given the high risk for hematologic relapse in patients with prior accelerated or blast phase, post-transplant TKI therapy should be considered for at least one year in this cohort of patients who are in remission following allogeneic HCT.³¹⁶⁻³¹⁸

TKI with or without DLI or omacetaxine can be considered for patients who are not in remission or in cytogenetic relapse or those with an increasing level of molecular relapse. The selection of TKI depends on prior TKI, the side effect profile of the TKI under consideration, the presence of comorbidities, and *BCR-ABL1* mutation status. Pre-existing mutations in the *BCR-ABL1* kinase domain, frequently associated with resistance to TKIs, are detectable in the majority of patients who relapse after allogeneic HCT.³¹⁹ *BCR-ABL1* mutational analysis is therefore essential prior to the selection of TKI for the treatment of post-transplant relapse.

In patients with CML that has previously failed imatinib, there are no data to support the use of post-transplant imatinib, and dasatinib, nilotinib, bosutinib, ponatinib, or omacetaxine may be more appropriate options. However, there are no data to support the use of post-transplant bosutinib,



NCCN Guidelines Version 2.2022

Chronic Myeloid Leukemia

ponatinib, or omacetaxine. CNS relapse of CML following allogeneic HCT has been described in few case reports.^{320,321} Dasatinib may also be an effective treatment for extramedullary relapse following allogeneic HCT.^{254,322,323} Participation in a clinical trial is highly desirable.

Emerging Treatment Options

Novel BCR-ABL1 inhibitors and small molecule inhibitors targeting several BCR-ABL1-independent pathways have demonstrated efficacy in preliminary clinical trials.^{324,325} These novel agents (either as monotherapy or in combination with currently approved TKIs) are being evaluated in ongoing clinical trials in all three phases of CML. Results from selected published clinical trials of novel agents are outlined in [Table 13](#).

The use of low-dose interferon in combination with TKI for a limited period prior to discontinuation of TKI, and gradual de-escalation of TKI prior to discontinuation of TKI therapy in patients with DMR are also being explored in ongoing clinical trials as potential strategies to improve TFR outcome.^{325,326} Pegylated interferon in combination with TKIs has demonstrated promising results, and ongoing clinical trials are evaluating the combination of second-generation TKIs with various pegylated interferons.³²⁷

Immunologic approaches such as the use of BCR-ABL1 immune peptides, immune checkpoint blockade, leukemia-associated antigens, and dendritic cell vaccines are also being evaluated to improve molecular response.³²⁴

Management of CML During Pregnancy and Breastfeeding

The median age of disease onset is 65 years, but CML occurs in all age groups. The EUTOS population-based registry has reported that

approximately 37% of patients at the time of diagnosis are of reproductive age.³²⁸ Clinical care teams should be prepared to address issues relating to fertility and pregnancy as well as counsel these patients about the potential risks and benefits of treatment discontinuation and possible resumption of TKI therapy should CML recur during pregnancy.

TKI Therapy and Conception

TKI therapy appears to affect some male hormones at least transiently, but does not appear to have a deleterious effect on male fertility. Furthermore, the miscarriage or fetal abnormality rate is not elevated in female partners of men on TKI therapy.³²⁹⁻³³³

The situation is more complex for women, as TKI therapy during pregnancy has been associated with both a higher rate of miscarriage and fetal abnormalities. Limited evidence from case reports on women with CML exposed to imatinib, dasatinib, or nilotinib during pregnancy indicates the need for close monitoring, a prolonged washout period prior to pregnancy, and prompt consideration of holding TKI therapy if pregnancy occurs while on imatinib, nilotinib, or dasatinib.³³⁴⁻³³⁸ In one report on the outcome of pregnancies in 180 women exposed to imatinib during pregnancy, 50% of pregnancies with known outcome were normal and 10% of pregnancies with known outcome had fetal abnormalities.³³⁴ Eighteen pregnancies ended in spontaneous abortion. In another report on the outcomes of pregnancy and conception during treatment with dasatinib, among 46 women treated with dasatinib, 15 women (33%) delivered a normal infant.³³⁵ Elective or spontaneous abortions were reported in 18 women (39%) and 8 women (17%), respectively, and 5 women (11%) had an abnormal pregnancy. Fetal abnormalities were reported in 7 cases. Among 33 women who conceived with dasatinib-treated men, 30 (91%) delivered infants who were normal at birth. Although there are no data regarding the outcome of pregnancy in



NCCN Guidelines Version 2.2022

Chronic Myeloid Leukemia

patients receiving bosutinib or ponatinib at the time of conception, these agents must be considered unsafe for use in pregnant women.

Discontinuation of TKI therapy because of pregnancy in women who were not in DMR ($\geq$ MR4.0; $\leq$ 0.01% *BCR-ABL1* IS) has only been reported in small series of patients.³³⁹⁻³⁴² In one series, among 10 women who stopped imatinib because of pregnancy after a median of 8 months of therapy, five of the nine women who had achieved a CHR lost the response after stopping therapy, and six had an increase in Ph-positive metaphases.³³⁹ At 18 months after resuming therapy, all nine patients had achieved a CHR but only three women achieved a CCyR and none had achieved an MMR. In another series that reported the outcomes of seven women who were not in DMR at the time imatinib was stopped because of pregnancy, three were in an MMR.³⁴⁰ All seven women had disease relapse. The three women who had an MMR at the time imatinib was stopped were able to regain the same response once the drug was restarted, whereas the remaining four patients were not.

Depending on other factors such as age, a natural pregnancy may occur months after stopping TKI therapy. Assuming the earliest time a woman could conceive and give birth naturally, without any washout period, is 10 months after stopping TKI, the likelihood is about 60% that her PCR will become positive if she was in DMR at the time of getting pregnant.^{339,340}

Conception while on active TKI therapy is strongly discouraged due to the risk of fetal abnormalities. Prior to attempting pregnancy, women and their partners should be counseled that no guidelines exist regarding how best to monitor CML during pregnancy, nor how best to manage progressive disease should it occur during pregnancy. Fertility preservation should be discussed with all patients of childbearing age

prior to the initiation of TKI therapy. Referral to a CML specialty center and consultation with a high-risk obstetrician is recommended.

Planning a Pregnancy

In men, the general recommendation is that TKI therapy need not be discontinued if a pregnancy is planned. However, experience is limited. Sperm banking can also be performed prior to starting TKI therapy, although there are no data regarding quality of sperm in men with untreated CML.

In women, due to the risk of miscarriage and fetal abnormalities during pregnancy, TKI therapy should be stopped prior to natural conception and the patient should remain off therapy during pregnancy.³³⁴⁻³³⁶ Referral to an in vitro fertilization (IVF) center is recommended in coordination with the patient's obstetrician. TKI should be stopped prior to attempting a natural pregnancy or oocyte retrieval, but the optimal timing of discontinuation is unknown. Compounding the high incidence of disease recurrence off TKI therapy are the significant obstacles that exist for women who choose one of the above forms of IVF, chief among which is the lack of access to centers that perform the procedure, high costs associated with the drugs and surgical procedures that may not be covered by insurance, costs of embryo/oocyte storage, and access to surrogate programs. Some women may require more than one IVF cycle to obtain enough potentially viable embryos for implantation. In addition, women may need a family medical leave from work to attend IVF appointments. It is also important to note that not all states allow surrogacy.

Treatment and Monitoring During Pregnancy

Most of the literature regarding treatment during pregnancy consists of case reports. The use of TKI therapy, particularly during the first trimester, should be avoided. If TKI therapy should be considered during



NCCN Guidelines Version 2.2022

Chronic Myeloid Leukemia

pregnancy, the potential benefit for the mother and the potential risk to the fetus of continuing TKI therapy versus the risk of treatment interruption leading to the loss of optimal disease response must be carefully evaluated on an individual basis prior to initiation of TKI therapy.

Interferon alpha and hydroxyurea have been used during pregnancy.³⁴³⁻³⁵¹ If treatment is deemed necessary during pregnancy, interferon can induce and maintain hematologic remission; if introduced earlier, interferon can preserve molecular remission after discontinuation of TKI.^{352,353} It is preferable to initiate treatment with interferon and the panel recommends against the use of hydroxyurea during pregnancy, especially in the first trimester, if possible. Data are insufficient to establish the use of peginterferon alfa-2a (risk category C) in pregnancy, and it should be used only if benefits outweigh potential risk to the fetus.³⁵⁴

Leukapheresis can be used for a rising white blood cell (WBC) count, although there are no data that recommend at what level of WBC count leukapheresis should be initiated.^{350,355-357} Low-dose aspirin or low-molecular-weight heparin can be considered for patients with thrombocytosis.^{358,359}

Monthly monitoring with qPCR and initiating treatment if the *BCR-ABL1* IS increases to >1.0% is recommended.

Breastfeeding

TKI therapy can be restarted after delivery. However, women on TKI therapy should be advised not to breastfeed, as TKIs pass into human breast milk.³⁶⁰⁻³⁶³

Breastfeeding without TKI treatment may be safe with molecular monitoring, but preferably in those patients with CML who have durable DMR. It may be acceptable to avoid TKIs for the short period of the first 2 to 5 days after labor to give the child colostrum.^{363,364}

Close molecular monitoring is recommended for women who extend the treatment-free period for breastfeeding. If the loss of MMR after treatment cessation is confirmed, breastfeeding needs to be terminated and TKI should be restarted.³⁶³

Specific Considerations for Children with CML

CML accounts for less than 3% of all pediatric leukemias. In general, children are diagnosed at a median age of 11 to 12 years, with approximately 10% presenting in advanced phase. Due to its rarity, there are no evidence-based recommendations for the management of CML in the pediatric population. Many pediatric oncologists follow treatment guidelines that are designed for adult patients. However, clinical presentations and host factors are different between children and adults, and several factors should be considered when treating pediatric patients with CML.³⁶⁵

Selection of TKI

Imatinib, dasatinib, and nilotinib are currently approved for treatment of CML in children.³⁶⁶⁻³⁶⁸ Higher dose imatinib (340 mg/m²) has also been shown to be effective and well tolerated in children.³⁶⁹⁻³⁷¹ There are very little data on the safety and efficacy of bosutinib and ponatinib in children.³⁷²

The validity of prognostic scores (eg, Sokal, Euro) for risk assessment or to make treatment decisions has not been established in the pediatric population.³⁷³ The ELTS score has demonstrated better differentiation of PFS than Sokal and Euro scores in children treated with imatinib.³⁷⁴

Monitoring for Long-Term Side Effects

Children have a much longer life expectancy than adults and TKI therapy may be needed for many decades; therefore, there are potential long-term side effects (such as delayed growth, changes in bone metabolism, thyroid



NCCN Guidelines Version 2.2022

Chronic Myeloid Leukemia

abnormalities, and effects on puberty and fertility) that may not be seen in adults.³⁷⁵

A number of studies have reported impaired longitudinal growth in children with CML treated with TKI therapy, and the effect is more significant when treatment was initiated during prepubertal age.³⁷⁶⁻³⁸² Growth should be monitored closely and a bone age x-ray should be obtained if longitudinal growth is delayed. A dual-energy x-ray absorptiometry (DEXA) scan should be obtained if bone mineral density is decreased on plain radiograph or if there is unprovoked fracture. Further evaluation and referral to an endocrinologist is also warranted.

In an international study (STOP IMAPED) that evaluated the discontinuation of imatinib in 14 children who were in sustained DMR, the overall probability of maintaining DMR at 6 months was 29%.³⁸³ Further studies in a larger cohort of patients are needed to identify the criteria for discontinuation of TKI therapy in the pediatric population. Therefore, discontinuation of TKI therapy in children at this time is not recommended outside the context of a clinical trial.³⁸⁴

Immunizations

There are little data on immune function of patients on TKI therapy, and TKI therapy potentially hinders routine vaccination for children with CML.³⁸⁵ In general, the use of inactivated killed vaccines for children on TKI therapy is safe, although it is unknown whether responses are comparable to those seen in healthy children. A study showed a higher seroconversion rate to H1N1 influenza vaccine in adult CML patients compared to patients with B-cell malignancies or HCT recipients.³⁸⁶

Administration of live vaccines during TKI therapy is not recommended in general, although one study showed that varicella vaccine could be safely given to some children with immune deficiency.³⁸⁷ Live attenuated annual influenza vaccine (nasal spray) should be avoided, and the inactivated

killed vaccine (flu shot) should be used for children receiving TKI therapy. Live vaccines could be considered after stopping TKI therapy for several weeks in patients with a DMR. In the United States, all required live vaccines are completed by the age of 4 to 6 years (<http://www.cdc.gov/vaccines/>). As CML is rarely seen in children younger than this age, few patients face this issue.



NCCN Guidelines Version 2.2022

Chronic Myeloid Leukemia

Table 1: First-Line TKI Therapy for CP-CML: Long-Term Follow-up Data from Phase III Studies

Trial	Study Arms	No. of Patients	Median Follow-up	CCyR ^a	MMR ^b	Disease Progression n (%)	PFS ^c	OS ^c
IRIS^{74, d}	Imatinib (400 mg once daily)	553	11 years	83%	—	38 (7%)	92%	83%
	Interferon alpha plus low-dose cytarabine	553		—	—	71 (13%)	—	79% ^e
DASISION⁷⁵	Dasatinib (100 mg once daily)	259	5 years	—	76% (<i>P</i> = .002)	12 (5%)	85%	91%
	Imatinib (400 mg once daily)	260		—	64%	19 (7%)	86%	90%
ENESTnd⁷⁶	Nilotinib (300 mg twice daily)	282	5 years	—	77% (<i>P</i> < .0001)	10 (4%)	92%	94%
	Imatinib (400 mg once daily)	283		—	60%	21 (7%)	91%	92%
BFORE^{77, f}	Bosutinib (400 mg once daily)	268	12 months	77% (<i>P</i> = .0075)	47% (<i>P</i> = .02)	4 (2%)	—	—
	Imatinib (400 mg once daily)	268		66%	37%	6 (3%)	—	—

CCyR, complete cytogenetic response; MMR, major molecular response ($\leq 0.1\%$ *BCR-ABL1* IS); OS, overall survival; PFS, progression-free survival

a. Primary endpoint of DASISION study: Confirmed CCyR rate at 12 months.

b. Primary endpoint of ENESTnd and BFORE studies: MMR ($\leq 0.1\%$ *BCR-ABL1* IS) rate at 12 months.

c. Long-term primary endpoint of IRIS trial in the imatinib group.

d. Due to the high rate of crossover to imatinib (66%) and the short duration of therapy (<1 year) before crossover among patients who had been randomly assigned to interferon alfa plus cytarabine, the long-term follow-up data focused on patients who had been randomly assigned to receive imatinib.

e. Data include survival among the 363 patients who crossed over to imatinib.

f. There were no differences in survival rates between the two treatment arms after a minimum follow-up of 12 months; long-term follow-up is ongoing.



NCCN Guidelines Version 2.2022

Chronic Myeloid Leukemia

Table 2: High-Dose Imatinib as First-Line Therapy for CP-CML: Long-Term Follow-up Data from Phase III Studies

Trial	Study Arms	No. of Patients	Median Follow-up	MMR	MR4.5	PFS	OS
TOPS study ^{78,a}	Imatinib (800 mg once daily)	319	42 months	79%	—	96% at 48 months	93% at 48 months
	Imatinib (400 mg once daily)	157		76%	—	94% at 48 months	94% at 48 months
CML IV study ^{80,b}	Imatinib (800 mg once daily)	420	10 years	89%	71%	77%	79%
	Imatinib (400 mg once daily)	400		92%	67%	80%	80%
SWOG study ^{79,c}	Imatinib (800 mg once daily)	73	12 months	53%	19%	92% (4-year PFS)	95% (4-year OS)
	Imatinib (400 mg once daily)	72		36%	9%	80% (4-year PFS)	90% (4-year OS)

MMR, major molecular response ($\leq 0.1\%$ *BCR-ABL1* IS); MR, molecular response;
MR4.5: ≥ 4.5 -log reduction in *BCR-ABL1* transcripts from baseline; OS, overall survival; PFS, progression-free survival

a. Primary endpoint: MMR rate at 12 months ($\leq 0.1\%$ *BCR-ABL1*), which corresponds to a 3-log reduction in *BCR-ABL1* transcripts compared with the standardized baseline established in IRIS study.

b. Primary endpoint: The impact of MMR on survival at 12 months. This study had 5 treatment arms (imatinib 400 mg once daily alone; imatinib 800 mg twice daily; imatinib 400 mg once daily with interferon or cytarabine; and imatinib after interferon failure). Only the data for imatinib 400 mg once daily alone vs. imatinib 800 mg twice daily are included in this table.

c. Primary endpoint: MR4.0 (≥ 4 -log reduction in *BCR-ABL1* transcripts from baseline) at 12 months. Results from the first part of SWOG S0325 study; follow-up after 12 months was not required for this study.



NCCN Guidelines Version 2.2022

Chronic Myeloid Leukemia

Table 3A: First-Line TKI Therapy for CP-CML: Molecular Response Rates According to Risk Score

Trial	Study Arms	Low-risk		Intermediate-risk		High-risk	
		MMR	MR4.5	MMR	MR4.5	MMR	MR4.5
DASISION ⁷⁵ (Euro risk score)	Dasatinib (100 mg once daily)	90%	55%	71%	43%	67%	31%
	Imatinib (400 mg once daily)	69%	44%	65%	28%	54%	30%
ENESTnd ⁷⁶ (Sokal risk score)	Nilotinib (300 mg twice daily)	—	53%	—	60%	—	45%
	Imatinib (400 mg once daily)	—	37%	—	33%	—	23%
BFORE ⁷⁷ (Sokal risk score)	Bosutinib (400 mg once daily)	58%	—	45%	—	34%	—
	Imatinib (400 mg once daily)	46%	—	39%	—	17%	—

Table 3B: First-Line TKI Therapy for CP-CML: Five-year Outcomes According to Sokal Risk Score

Trial	Study Arms	Low-risk			Intermediate-risk			High-risk		
		Disease Progression n (%)	PFS	OS	Disease Progression n (%)	PFS	OS	Disease Progression n (%)	PFS	OS
ENESTnd ⁷⁶ (Sokal risk score)	Nilotinib (300 mg twice daily)	1 (1%)	96%	97%	2 (2%)	93%	94%	7 (9%)	86%	89%
	Imatinib (400 mg once daily)	0%	100%	100%	10 (10%)	88%	89%	11 (14%)	83%	84%

MMR, major molecular response ($\leq 0.1\%$ *BCR-ABL1* IS); MR, molecular response; MR4.5: 4.5-log reduction in *BCR-ABL1* transcripts from baseline; OS, overall survival; PFS, progression-free survival



NCCN Guidelines Version 2.2022

Chronic Myeloid Leukemia

Table 4. Adverse Events of First-Line TKI Therapy in CP-CML

Toxicity	DASISION ⁷⁵		ENESTnd ⁷⁶		BFORE ⁷⁷	
	Dasatinib 100 mg QD	Imatinib 400 mg QD	Nilotinib 300 mg BID	Imatinib 400 mg QD	Bosutinib 400 mg QD	Imatinib 400 mg QD
Hematologic toxicities (Grade 3/4)						
Anemia	13%	9%	4%	6%	3%	5%
Neutropenia	29%	24%	12%	22%	7%	12%
Thrombocytopenia	22%	14%	10%	9%	14%	6%
Biochemical abnormalities (Grade 3/4)						
Increased lipase	NR	NR	9%	4%	13%	6%
Increased glucose	NR	NR	7%	<1%	2%	2%
Decreased phosphate	7%	28%	8%	10%	5%	17%
Increased ALT	NR	NR	4%	2%	23%	3%
Increased AST	NR	NR	NR	NR	12%	3%
Nonhematologic toxicities (any grade) *						
Rash	13%	18%	38%	19%	20%	13%
Headache	13%	11%	32%	23%	19%	13%
Fatigue	9%	11%	23%	20%	19%	18%
Muscle spasms	23%	41%	12%	34%	2%	26%
Peripheral edema	13%	37%	9%	20%	4%	14%
Pleural effusion	28%	<1%	2%	1%	NR	NR
Hypertension	NR	NR	10%	4%	NR	NR
Pulmonary hypertension	5%	<1%	0%	0%	NR	NR
Diarrhea	21%	22%	19%	46%	70%	34%
Constipation	NR	NR	20%	8%	NR	NR
Nausea	10%	24%	22%	41%	35%	39%
Vomiting	5%	11%	15%	27%	18%	16%

ALT, alanine aminotransferase; AST, aspartate aminotransferase; BID, twice daily; QD, once daily.

* Non-hematologic toxicities from the DASISION study (except pleural effusion) are from the 3-year follow-up. No new adverse events were observed with 5-year follow-up.



NCCN Guidelines Version 2.2022

Chronic Myeloid Leukemia

Table 5. Early Molecular Response ($\leq 10\%$ *BCR-ABL1* IS at 3 months) After First-Line TKI Therapy and Survival Outcomes

Trial	Study Arms	5-year PFS		5-year OS	
		<i>BCR-ABL1</i> $\leq 10\%$	<i>BCR-ABL1</i> $>10\%$	<i>BCR-ABL1</i> $\leq 10\%$	<i>BCR-ABL1</i> $>10\%$
DASISION⁷⁵	Dasatinib (100 mg once daily)	89%	72%	94%	81%
	Imatinib (400 mg once daily)	93%	72%	95%	81%
ENESTnd⁷⁶	Nilotinib (300 mg twice daily)	95%	78%	98%	82%
	Nilotinib (400 mg twice daily)	96%	89%	96%	93%
	Imatinib (400 mg once daily)	98%	79%	99%	79%
CML IV Study¹¹⁸	Imatinib (400 mg once daily)	92%	87%	94%	87%

OS, overall survival; PFS, progression-free survival

progress



NCCN Guidelines Version 2.2022

Chronic Myeloid Leukemia

Table 6. Second-Line and Subsequent TKI Therapy for CP-CML: Long-Term Follow-up Data from Phase II/III Studies

TKI	No. of Patients	Median Follow-up	MCyR	CCyR	MMR	PFS	OS
Dasatinib^{137,a} (100 mg once daily)	Imatinib-R (n = 124)	7 years	—	—	43%	39%	63%
	Imatinib-I (n = 43)		—	—	55%	51%	70%
Nilotinib^{138,b} (400 mg twice daily)	Imatinib-R (n = 226)	4 years	59%	45%	—	57%	78%
	Imatinib-I (n = 95)						
Bosutinib^{147,k} (400 mg once daily)	Imatinib and dasatinib-R (n = 38)	4 years	39%	22%	—	—	67%
	Imatinib and dasatinib-I (n = 50)		42%	40%	—	—	80%
	Imatinib and nilotinib-R (n = 26)		38%	31%	—	—	87%
Ponatinib^{140,c} (45 mg once daily)	Dasatinib or nilotinib-R or I (n = 203)	57 months	56%	49%	35%	52% at 5 years	76% at 5 years
	T315I mutation (n = 64)		72%	70%	58%	50% at 5 years	66% at 5 years

R = Resistant; I = Intolerant; CCyR, complete cytogenetic response; MCyR, major cytogenetic response; MMR, major molecular response ($\leq 0.1\%$ *BCR-ABL1* IS); OS, overall survival; PFS, progression-free survival

a. Primary endpoint: MCyR rate at 6 months when administered 100 mg once daily vs. 70 mg twice daily.

b. Primary endpoint: MCyR rate in patients with imatinib intolerance or imatinib-resistant disease.

c. Primary endpoint: MCyR at any time within the first 12 months.



NCCN Guidelines Version 2.2022

Chronic Myeloid Leukemia

Table 7. Early Molecular Response ($\leq 10\%$ *BCR-ABL1* IS) After Second-Line TKI Therapy and Survival Outcomes

TKI	Median Follow-up	Progression-Free Survival (PFS)				Overall Survival (OS)			
		<i>BCR-ABL1</i> $\leq 10\%$		<i>BCR-ABL1</i> $>10\%$		<i>BCR-ABL1</i> $\leq 10\%$		<i>BCR-ABL1</i> $>10\%$	
		3 months	6 months	3 months	6 months	3 months	6 months	3 months	6 months
Dasatinib¹³⁷ (100 mg once daily)	7 years	56%	57%	21%	4%	72%	74%	56%	50%
Nilotinib¹³⁸ (400 mg twice daily)	4 years	67%	58%	42%	39%	81%	82%	71%	73%

progress



NCCN Guidelines Version 2.2022

Chronic Myeloid Leukemia

Table 8. Responses Based on BCR-ABL1 Mutation Status^a

Mutation	Major Cytogenetic Response (MCyR), n/N (%)			
	Bosutinib ¹⁴⁷	Dasatinib ¹⁹¹	Nilotinib ¹⁹²	Ponatinib ¹⁹⁷
E255K ^b	—	9/16 (56%)	3/7 (43%)	8/13 (62%)
E255V ^b	—	4/11 (36%)		1/4 (25%)
E459K ^b	—	—	—	3/7 (43%)
F317L ^{c,d}	1/7 (14%)	2/14 (14%)	—	13/29 (45%)
F359C ^b	1/2 (50%)	3/5 (60%)	1/11 (9%)	1/7 (14%)
F359V ^b	2/3 (67%)	17/27 (63%)		11/20 (55%)
F359I ^b	2/2 (100%)	10/12 (83%)	—	3/4 (75%)
G250E ^{b,e}	0/5 (0%)	29/60 (48%)	3/5 (60%)	8/12 (67%)
H396R	—	17/33 (52%)	—	1/5 (20%)
L248V	—	10/15 (67%)	—	1/2 (50%)
M244V	2/3 (67%)	27/26 (59%)	—	4/9 (56%)
M351T	—	28/54 (52%)	—	1/2 (50%)
Y253H ^b	5/6 (83%)	15/23 (65%)	1/8 (13%)	1/2 (50%)
V299L ^{c,e}	0/2 (0%)	—	—	3/8 (38%)

a. Mutations contraindicated for imatinib are too numerous to include. There are compound mutations that can cause resistance to ponatinib, but those are uncommon following treatment with bosutinib, dasatinib, or nilotinib.

b. *BCR-ABL1* mutations resistant to nilotinib.

c. *BCR-ABL1* mutations resistant to dasatinib.

d. Bosutinib has minimal activity against F317L mutation and in vitro studies suggest that F317L is highly sensitive to nilotinib.

e. *BCR-ABL1* mutations resistant to bosutinib.



NCCN Guidelines Version 2.2022

Chronic Myeloid Leukemia

Table 9. Adverse Events of Second-Line and Subsequent TKI Therapy in CP-CML

Toxicity (any grade)	Dasatinib ¹³⁷ (100 mg QD)	Nilotinib ¹³⁸ (300 mg BID)	Bosutinib ¹⁴⁷ (400 mg QD)	Ponatinib ¹⁴⁰ (45 mg QD)
Rash	33%	31%	28%	47%
Headache	—	18%	27%	43%
Fatigue	37%	21%	24%	30%
Myalgias/Arthralgias	38%	11%	18%	24%/33%
Pleural effusion	28%	—	17%	—
Hypertension	—	—	8%	37%
Hemorrhage	26%	—	—	—
Diarrhea	42%	12%	83%	20%
Constipation	—	13%	13%	41%
Nausea	27%	25%	48%	29%
Vomiting		13%	38%	19%
Increased blood creatinine	—	—	13%	—
Increased lipase	—	—	—	27%
Increased ALT/AST	—	—	15%	—

ALT, alanine aminotransferase; AST, aspartate aminotransferase; BID, twice daily; QD, once daily.



NCCN Guidelines Version 2.2022

Chronic Myeloid Leukemia

Table 10. Summary of Limited Longer-Term Follow-up Data from the TKI Discontinuation Trials

Trial	Treatment Prior to Discontinuation	No. of Patients	Depth and Duration of MR Required for Discontinuation	Trigger to Resume TKI Therapy	Median Follow-up	Treatment-free Remission (TFR) Rate
STIM1 ²¹³	Imatinib ± interferon	100	MR5.0 for at least 2 years	Loss of MR5.0	77 months	38% at 60 months
TWISTER ²¹⁸	Imatinib ± interferon	40	MR4.5 for at least 2 years	Loss of MR5.0	103 months	45% (molecular relapse-free survival 45% at 8 years)
HOVON ²¹⁴	Imatinib + cytarabine	15	MR4.5 for at least 2 years	Loss of MR4.5	36 months	33% at 24 months
A-STIM ²¹⁵	Imatinib ± interferon	80	MR5.0 for at least 2 years	Loss of MMR	31 months	61% at 36 months
ISAV study ²¹⁶	Imatinib (after failure of interferon or hydroxyurea)	108	CMR for at least 18 months	Loss of MMR	36 months	52% at 36 months
KID study ²¹⁷	Imatinib ± interferon	90	MR4.5 for at least 2 years	Loss of MMR	27 months	59% at 24 months
Stop 2G-TKI ²¹⁹	Dasatinib/Nilotinib (first- or second-line)	60	MR4.5 for at least 24 months	Loss of MMR	47 months	54% at 48 months
DASFREE ²²⁴	Dasatinib (first- or second-line)	84	MR4.5 for 12 months	Loss of MMR	2 years	46% at 24 months
ENESTFreedom ²²⁰	Nilotinib (first-line)	190	MR4.5 for 12 months	Loss of MMR	96 weeks	49% at 96 weeks
ENESTop study ²²¹	Nilotinib (second-line)	126	MR4.5 for 12 months	Loss of MMR	96 weeks	53% at 96 weeks
DADI ²²⁵	Dasatinib (first-line)	68	MR4.5 for at least 24 months	Loss of MMR	23 months	55% at 6 months
DADI ²²²	Dasatinib (second-line)	63	MR4.0 for at least 12 months	Loss of MR4.0	44 months	44% at 36 months
EURO-SKI ²²³	Any TKI	758	MR4.0 for at least 1 year	Loss of MMR	27 months	50% at 24 months

CMR, complete molecular response (undetectable *BCR-ABL1* by qPCR as determined by local laboratories); MMR, major molecular response ($\leq 0.1\%$ *BCR-ABL1* IS); MR, molecular response; MR4.0, $\leq 0.01\%$ *BCR-ABL1* IS; MR4.5, $\leq 0.0032\%$ *BCR-ABL1* IS or >4.5 -log reduction of *BCR-ABL1* and undetectable minimal residual disease on qPCR with a sensitivity of ≥ 4.5 -log reduction; MR5.0, 5-log reduction in *BCR ABL1* levels and undetectable minimal residual disease on qPCR with a sensitivity of ≥ 4.5 -log reduction.



NCCN Guidelines Version 2.2022

Chronic Myeloid Leukemia

Table 11. TKI Therapy for Disease Progression to AP-CML: Long-Term Follow-up Data from Phase II/III Studies

TKI	No. of Patients	Median Follow-up	MCyR	CCyR	OS	PFS
Dasatinib^{233,a} (140 mg once daily)	Imatinib-R (n = 117)	24 months	36%	29%	63%	51%
	Imatinib-I (n = 41)		46%	41%		
Nilotinib^{236,b} (400 mg twice daily)	Imatinib-R (n = 109)	24 months	30%	19%	70%	33%
	Imatinib-I (n = 27)		41%	30%		
Bosutinib^{238,c} (500 mg once daily)	Prior imatinib only (n = 49)	48 months	48%	35%	66%	—
	Imatinib followed by dasatinib or nilotinib (n = 30)		27%	23%	45%	—
Ponatinib^{140,d} (45 mg once daily)	Dasatinib or nilotinib-R or I (n = 65)	32 months	45%	28%	48% at 5 years	19% at 5 years
	T315I mutation (n = 18)		67%	44%	52% at 5 years	29% at 5 years

R = Resistant; I = Intolerant; CCyR, Complete cytogenetic response; MCyR, major cytogenetic response; OS, overall survival; PFS, progression-free survival

a. Primary endpoint: Major hematologic response (MaHR). The rate of MaHR at 5 years was 67% for 140 mg once daily and 69% for 70 mg twice daily (Ottmann O, et al. Blood Cancer J 2018;8:88).

b. Primary endpoint: Confirmed complete hematologic response rate, achieved in 30% of patients with imatinib-resistant disease and 37% of imatinib-intolerant patients.

c. Primary endpoint: Confirmed overall hematologic response by 48 weeks.

d. Primary endpoint: MaHR at any time within the first 6 months.



NCCN Guidelines Version 2.2022

Chronic Myeloid Leukemia

Table 12. TKI Therapy for Disease Progression to BP-CML: Long-term Follow-up Data from Phase II/III Studies

TKI	No. of Patients	Median Follow-up	MCyR	CCyR	OS
Dasatinib ^{a,235} (140 mg once daily)	Lymphoid blast phase (n = 33)	24 months	50%	38%	21%
	Myeloid blast phase (n = 75)		25%	14%	24%
Nilotinib ^{b,237} (400 mg twice daily)	Lymphoid blast phase (n = 31)	24 months	52%	32%	10%
	Myeloid blast phase (n = 105)		38%	30%	32%
Bosutinib ^{c,238} (500 mg once daily)	Prior imatinib only (n = 36)	48 months	50%	37%	28%
	Imatinib followed by dasatinib or nilotinib (n = 28)		21%	17%	17%
Ponatinib ^{d,140} (45 mg once daily)	Dasatinib or nilotinib-R or I (n = 38)	6 months	18%	16%	9% at 3 years
	T315I mutation (n = 24)		29%	21%	

R = Resistant; I= Intolerant; CCyR, complete cytogenetic response; MCyR, major cytogenetic response; OS, overall survival

a. Primary endpoint: Major hematologic response (MaHR).

b. Endpoints: Duration of MaHR, MCyR, and OS.

c. Primary endpoint: Confirmed overall hematologic response by 48 weeks.

d. MaHR at any time within the first 6 months.



NCCN Guidelines Version 2.2022

Chronic Myeloid Leukemia

Table 13: Results from Selected Published Clinical Trials Evaluating Novel Treatment Options

Drug Class	Clinical Trial	TKI	No. of Patients	Median Follow-up	Response Rates
BCR-ABL1 inhibitors	Phase I (Dose-escalation study) ³⁸⁸ CP-CML or AP-CML with resistance or intolerance to at least 2 previous TKIs	Asciminib (10–200 mg once or twice daily)	CP-CML without T315I (n = 113)	72 weeks	MCyR: 77%; CCyR: 70%
			CP-CML with T315I (n = 28)	37 weeks	MCyR: 60%; CCyR: 44%
			AP-CML without T315I (n = 4)	46 weeks	CHR: 100%; CCyR: 0%
			AP-CML with T315I (n = 5)	16 weeks	CHR: 80%; CCyR: 20%
	Phase III (REPRISE study) ³⁸⁹ Newly diagnosed CP-CML	Radotinib (300 mg twice daily)	n = 79	≥48 months	MMR: 85%; MR4.5: 58%
		Radotinib (400 mg twice daily)	n = 81		MMR: 83%; MR4.5: 56%
		Imatinib (400 mg once daily)	n = 81		MMR: 75%; MR4.5: 49%
	Phase II ³⁹⁰ CP-CML or AP-CML with resistance or intolerance to imatinib	Radotinib (400 mg twice daily)	n = 77	23 months	MCyR: 65%; CCyR: 47%; MMR: 14%
Aurora kinase inhibitors	Phase I ³⁹¹ CP-CML or AP/BP-CML after failure of imatinib	Lonafarnib (100 mg twice daily) + imatinib (400 mg once daily)	CP-CML (n = 9)		CHR: 9%; CCyR: 4%
		Lonafarnib (100 mg twice daily) + imatinib (600 mg once daily)	AP/BP-CML (n = 14)		CHR: 14%; PCyR: 4%
	Phase I ³⁹² CP-CML CML after failure of imatinib	Tipifarnib (300 mg twice daily) + imatinib (400 mg once daily)	n = 26		CHR: 68%; CCyR: 12%
Farnesyl transferase inhibitors	Phase I (Dose-escalation study) ³⁹³ AP-CML or BP-CML with resistance or intolerance to previous TKIs	Danuserib (180 mg/m ² ; 3-hour IV infusion; days 1–7; 14-day cycle)	AP-CML (n = 7)		CHR: 3%
			BP-CML (n = 9)		CCyR: 4%
	Phase II ³⁹⁴ CP-CML, AP/BP-CML with T315I mutation	Tozasertib (5-day continuous IV infusion every 14 days at 40 mg/m ² /h, 32 mg/m ² /h, or 24 mg/m ² /h)	CP-CML (n = 15)		CHR: 7%; MCyR: 13%; CCyR: 13%
			AP-CML (n = 14)		MCyR: 7%; CCyR: 7%
			BP-CML (n = 11)		MCyR: 9%
JAK2 inhibitors	Phase I (Dose-escalation study) ³⁹⁵ CP-CML with no history of disease progression to AP-CML or BP-CML	Ruxolitinib (5, 10, and 15 mg) + nilotinib (300 mg or 400 mg twice daily)	n = 11		40% had values between MMR and MR4.0; 10% had values between MR4.0 and MR4.5; and 40% had MR4.5



NCCN Guidelines Version 2.2022

Chronic Myeloid Leukemia

References

1. Siegel RL, Miller KD, Jemal A. Cancer statistics, 2020. CA Cancer J Clin 2020;70:7-30. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/31912902>.
2. Faderl S, Talpaz M, Estrov Z, et al. The biology of chronic myeloid leukemia. N Engl J Med 1999;341:164-172. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/10403855>.
3. Melo JV. The diversity of BCR-ABL fusion proteins and their relationship to leukemia phenotype. Blood 1996;88:2375-2384. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/8839828>.
4. Melo JV. BCR-ABL gene variants. Baillieres Clin Haematol 1997;10:203-222. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/9376660>.
5. Sawyers CL. Chronic myeloid leukemia. N Engl J Med 1999;340:1330-1340. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/10219069>.
6. Radich JP, Dai H, Mao M, et al. Gene expression changes associated with progression and response in chronic myeloid leukemia. Proc Natl Acad Sci U S A 2006;103:2794-2799. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/16477019>.
7. Jamieson CHM, Ailles LE, Dylla SJ, et al. Granulocyte-macrophage progenitors as candidate leukemic stem cells in blast-crisis CML. N Engl J Med 2004;351:657-667. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/15306667>.
8. U.S. National Library of Medicine Key MEDLINE® Indicators Available at: http://www.nlm.nih.gov/bsd/bsd_key.html.
9. Mitelman F. The cytogenetic scenario of chronic myeloid leukemia. Leuk Lymphoma 1993;11 Suppl 1:11-15. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/8251885>.
10. Douet-Guilbert N, Morel F, Le Charpentier T, et al. Interphase FISH for follow-up of Philadelphia chromosome-positive chronic myeloid leukemia treatment. Anticancer Res 2004;24:2535-2539. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/15330210>.
11. Seong DC, Kantarjian HM, Ro JY, et al. Hypermetaphase fluorescence in situ hybridization for quantitative monitoring of Philadelphia chromosome-positive cells in patients with chronic myelogenous leukemia during treatment. Blood 1995;86:2343-2349. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/7662980>.
12. Dewald GW, Wyatt WA, Juneau AL, et al. Highly sensitive fluorescence in situ hybridization method to detect double BCR/ABL fusion and monitor response to therapy in chronic myeloid leukemia. Blood 1998;91:3357-3365. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/9558393>.
13. Kantarjian HM, Talpaz M, Cortes J, et al. Quantitative polymerase chain reaction monitoring of BCR-ABL during therapy with imatinib mesylate (STI571; gleevec) in chronic-phase chronic myelogenous leukemia. Clin Cancer Res 2003;9:160-166. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/12538464>.
14. Hughes T, Deininger M, Hochhaus A, et al. Monitoring CML patients responding to treatment with tyrosine kinase inhibitors: review and recommendations for harmonizing current methodology for detecting BCR-ABL transcripts and kinase domain mutations and for expressing results. Blood 2006;108:28-37. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/16522812>.
15. Biernaux C, Loos M, Sels A, et al. Detection of major bcr-abl gene expression at a very low level in blood cells of some healthy individuals. Blood 1995;86:3118-3122. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/7579406>.
16. Bose S, Deininger M, Gora-Tybor J, et al. The presence of typical and atypical BCR-ABL fusion genes in leukocytes of normal individuals: biologic significance and implications for the assessment of minimal residual disease. Blood 1998;92:3362-3367. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/9787174>.
17. Baccarani M, Castagnetti F, Gugliotta G, et al. The proportion of different BCR-ABL1 transcript types in chronic myeloid leukemia. An international overview. Leukemia 2019;33:1173-1183. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/30675008>.



NCCN Guidelines Version 2.2022

Chronic Myeloid Leukemia

18. Hanfstein B, Lauseker M, Hehlmann R, et al. Distinct characteristics of e13a2 versus e14a2 BCR-ABL1 driven chronic myeloid leukemia under first-line therapy with imatinib. *Haematologica* 2014;99:1441-1447. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/24837466>.
19. Jain P, Kantarjian H, Patel KP, et al. Impact of BCR-ABL transcript type on outcome in patients with chronic-phase CML treated with tyrosine kinase inhibitors. *Blood* 2016;127:1269-1275. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/26729897>.
20. Castagnetti F, Gugliotta G, Breccia M, et al. The BCR-ABL1 transcript type influences response and outcome in Philadelphia chromosome-positive chronic myeloid leukemia patients treated frontline with imatinib. *Am J Hematol* 2017;92:797-805. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/28466557>.
21. Ercaliskan A, Eskazan AE. The impact of BCR-ABL1 transcript type on tyrosine kinase inhibitor responses and outcomes in patients with chronic myeloid leukemia. *Cancer* 2018;124:3806-3818. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/29694669>.
22. Pffirmann M, Evtimova D, Saussele S, et al. No influence of BCR-ABL1 transcript types e13a2 and e14a2 on long-term survival: results in 1494 patients with chronic myeloid leukemia treated with imatinib. *J Cancer Res Clin Oncol* 2017;143:843-850. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/28083711>.
23. Verma D, Kantarjian HM, Jones D, et al. Chronic myeloid leukemia (CML) with P190 BCR-ABL: analysis of characteristics, outcomes, and prognostic significance. *Blood* 2009;114:2232-2235. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/19531657>.
24. Arun AK, Senthamizhselvi A, Mani S, et al. Frequency of rare BCR-ABL1 fusion transcripts in chronic myeloid leukemia patients. *Int J Lab Hematol* 2017;39:235-242. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/28035733>.
25. Gong Z, Medeiros LJ, Cortes JE, et al. Clinical and prognostic significance of e1a2 BCR-ABL1 transcript subtype in chronic myeloid leukemia. *Blood Cancer J* 2017;7:e583. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/28708130>.
26. Qin YZ, Jiang Q, Jiang H, et al. Prevalence and outcomes of uncommon BCR-ABL1 fusion transcripts in patients with chronic myeloid leukaemia: data from a single centre. *Br J Haematol* 2018;182:693-700. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/29974949>.
27. Xue M, Wang Q, Huo L, et al. Clinical characteristics and prognostic significance of chronic myeloid leukemia with rare BCR-ABL1 transcripts. *Leuk Lymphoma* 2019;60:3051-3057. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/31258010>.
28. Verstovsek S, Lin H, Kantarjian H, et al. Neutrophilic-chronic myeloid leukemia: low levels of p230 BCR/ABL mRNA and undetectable BCR/ABL protein may predict an indolent course. *Cancer* 2002;94:2416-2425. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/12015767>.
29. Langabeer SE, McCarron SL, Kelly J, et al. Chronic Myeloid Leukemia with e19a2 BCR-ABL1 Transcripts and Marked Thrombocytosis: The Role of Molecular Monitoring. *Case Rep Hematol* 2012;2012:458716. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/22937329>.
30. Crampe M, Haslam K, Kelly J, et al. Characterization of a novel variant BCR-ABL1 fusion transcript in a patient with chronic myeloid leukemia: Implications for molecular monitoring. *Hematol Oncol Stem Cell Ther* 2017;10:85-88. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/27013275>.
31. Langabeer SE. Standardized Molecular Monitoring for Variant BCR-ABL1 Transcripts in Chronic Myeloid Leukemia. *Arch Pathol Lab Med* 2015;139:969. Available at:
32. Shanmuganathan N, Hughes TP. Molecular monitoring in CML: how deep? How often? How should it influence therapy? *Hematology Am Soc Hematol Educ Program* 2018;2018:168-176. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/30504306>.
33. Burmeister T, Reinhardt R. A multiplex PCR for improved detection of typical and atypical BCR-ABL fusion transcripts. *Leuk Res* 2008;32:579-585. Available at:
34. Bennour A, Ouahchi I, Moez M, et al. Comprehensive analysis of BCR/ABL variants in chronic myeloid leukemia patients using multiplex



NCCN Guidelines Version 2.2022

Chronic Myeloid Leukemia

RT-PCR. Clin Lab 2012;58:433-439. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/22783572>.

35. Mir R, Ahmad I, Javid J, et al. Simple multiplex RT-PCR for identifying common fusion BCR-ABL transcript types and evaluation of molecular response of the a2b2 and a2b3 transcripts to Imatinib resistance in north Indian chronic myeloid leukemia patients. Indian J Cancer 2015;52:314-318. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/26905124>.

36. Pagani IS, Dang P, Saunders VA, et al. Clinical utility of genomic DNA Q-PCR for the monitoring of a patient with atypical e19a2 BCR-ABL1 transcripts in chronic myeloid leukemia. Leuk Lymphoma 2020;1-3. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/32508223>.

37. Petiti J, Lo Iacono M, Dragani M, et al. Novel Multiplex Droplet Digital PCR Assays to Monitor Minimal Residual Disease in Chronic Myeloid Leukemia Patients Showing Atypical BCR-ABL1 Transcripts. J Clin Med 2020;9. Available at:

38. Cortes JE, Talpaz M, Giles F, et al. Prognostic significance of cytogenetic clonal evolution in patients with chronic myelogenous leukemia on imatinib mesylate therapy. Blood 2003;101:3794-3800. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/12560227>.

39. O'Dwyer ME, Mauro MJ, Blasdel C, et al. Clonal evolution and lack of cytogenetic response are adverse prognostic factors for hematologic relapse of chronic phase CML patients treated with imatinib mesylate. Blood 2004;103:451-455. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/14512312>.

40. Wang W, Cortes JE, Lin P, et al. Clinical and prognostic significance of 3q26.2 and other chromosome 3 abnormalities in CML in the era of tyrosine kinase inhibitors. Blood 2015;126:1699-1706. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/26243778>.

41. Wang W, Tang G, Cortes JE, et al. Chromosomal rearrangement involving 11q23 locus in chronic myelogenous leukemia: a rare phenomenon frequently associated with disease progression and poor prognosis. J Hematol Oncol 2015;8:32. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/25888368>.

42. Wang W, Cortes JE, Tang G, et al. Risk stratification of chromosomal abnormalities in chronic myelogenous leukemia in the era of tyrosine kinase inhibitor therapy. Blood 2016;127:2742-2750. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/27006386>.

43. Verma D, Kantarjian H, Shan J, et al. Survival outcomes for clonal evolution in chronic myeloid leukemia patients on second generation tyrosine kinase inhibitor therapy. Cancer 2010;116:2673-2681. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/20499401>.

44. Fabarius A, Kalmanti L, Dietz CT, et al. Impact of unbalanced minor route versus major route karyotypes at diagnosis on prognosis of CML. Ann Hematol 2015;94:2015-2024. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/26385387>.

45. Fabarius A, Leitner A, Hochhaus A, et al. Impact of additional cytogenetic aberrations at diagnosis on prognosis of CML: long-term observation of 1151 patients from the randomized CML Study IV. Blood 2011;118:6760-6768. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/22039253>.

46. Alhurairi A, Kantarjian H, Boddur P, et al. Prognostic significance of additional chromosomal abnormalities at the time of diagnosis in patients with chronic myeloid leukemia treated with frontline tyrosine kinase inhibitors. Am J Hematol 2018;93:84-90. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/29027261>.

47. Bumm T, Muller C, Al-Ali H-K, et al. Emergence of clonal cytogenetic abnormalities in Ph- cells in some CML patients in cytogenetic remission to imatinib but restoration of polyclonal hematopoiesis in the majority. Blood 2003;101:1941-1949. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/12411298>.

48. Feldman E, Najfeld V, Schuster M, et al. The emergence of Ph-, trisomy -8+ cells in patients with chronic myeloid leukemia treated with imatinib mesylate. Exp Hematol 2003;31:702-707. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/12901975>.

49. Medina J, Kantarjian H, Talpaz M, et al. Chromosomal abnormalities in Philadelphia chromosome-negative metaphases appearing during imatinib mesylate therapy in patients with Philadelphia chromosome-positive



NCCN Guidelines Version 2.2022

Chronic Myeloid Leukemia

chronic myelogenous leukemia in chronic phase. Cancer 2003;98:1905-1911. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/14584073>.

50. Terre C, Eclache V, Rousselot P, et al. Report of 34 patients with clonal chromosomal abnormalities in Philadelphia-negative cells during imatinib treatment of Philadelphia-positive chronic myeloid leukemia. Leukemia 2004;18:1340-1346. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/15190256>.

51. Deininger MW, Cortes J, Paquette R, et al. The prognosis for patients with chronic myeloid leukemia who have clonal cytogenetic abnormalities in philadelphia chromosome-negative cells. Cancer 2007;110:1509-1519. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/17702093>.

52. Jabbour E, Kantarjian HM, Abruzzo LV, et al. Chromosomal abnormalities in Philadelphia chromosome negative metaphases appearing during imatinib mesylate therapy in patients with newly diagnosed chronic myeloid leukemia in chronic phase. Blood 2007;110:2991-2995. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/17625066>.

53. Vignetti M, Fazi P, Cimino G, et al. Imatinib plus steroids induces complete remissions and prolonged survival in elderly Philadelphia chromosome-positive patients with acute lymphoblastic leukemia without additional chemotherapy: results of the Gruppo Italiano Malattie Ematologiche dell'Adulto (GIMEMA) LAL0201-B protocol. Blood 2007;109:3676-3678. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/17213285>.

54. Fabarius A, Haferlach C, Muller MC, et al. Dynamics of cytogenetic aberrations in Philadelphia chromosome positive and negative hematopoiesis during dasatinib therapy of chronic myeloid leukemia patients after imatinib failure. Haematologica 2007;92:834-837. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/17550857>.

55. Baldazzi C, Luatti S, Marzocchi G, et al. Emergence of clonal chromosomal abnormalities in Philadelphia negative hematopoiesis in chronic myeloid leukemia patients treated with nilotinib after failure of imatinib therapy. Leuk Res 2009;33:e218-220. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/19525008>.

56. Wang H, Jin J, Wang Y, et al. Clonal chromosomal abnormalities in Philadelphia-negative cells in chronic myeloid leukemia patients treated with nilotinib used in first-line therapy. Ann Hematol 2013;92:1625-1632. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/23793947>.

57. Ni H, Sun X, Xu Y, et al. Clinical implications of clonal chromosomal abnormalities in Philadelphia negative cells in CML patients after treated with tyrosine kinase inhibitors. Cancer Genet 2019;238:44-49. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/31425925>.

58. Sheng G, Xue M, Wang Q, et al. Occurrence of chromosomal abnormalities in Philadelphia chromosome-negative metaphases in patients with chronic-phase chronic myeloid leukemia undergoing TKI treatments. Leuk Lymphoma 2019;60:3503-3511. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/31282805>.

59. Issa GC, Kantarjian HM, Gonzalez GN, et al. Clonal chromosomal abnormalities appearing in Philadelphia chromosome-negative metaphases during CML treatment. Blood 2017;130:2084. Available at: <http://www.bloodjournal.org/content/130/19/2084.abstract>.

60. Karimata K, Masuko M, Ushiki T, et al. Myelodysplastic syndrome with Ph negative monosomy 7 chromosome following transient bone marrow dysplasia during imatinib treatment for chronic myeloid leukemia. Intern Med 2011;50:481-485. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/21372464>.

61. Navarro JT, Feliu E, Grau J, et al. Monosomy 7 with severe myelodysplasia developing during imatinib treatment of Philadelphia-positive chronic myeloid leukemia: two cases with a different outcome. Am J Hematol 2007;82:849-851. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/17563075>.

62. Bidet A, Dulucq S, Smol T, et al. Poor prognosis of chromosome 7 clonal aberrations in Philadelphia-negative metaphases and relevance of potential underlying myelodysplastic features in chronic myeloid leukemia. Haematologica 2019;104:1150-1155. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/30573507>.

63. Soverini S, De Benedittis C, Castagnetti F, et al. In chronic myeloid leukemia patients on second-line tyrosine kinase inhibitor therapy, deep sequencing of BCR-ABL1 at the time of warning may allow sensitive



NCCN Guidelines Version 2.2022

Chronic Myeloid Leukemia

detection of emerging drug-resistant mutants. BMC Cancer 2016;16:572. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/27485109>.

64. Kizilers A, Crisa E, Lea N, et al. Effect of low-level BCR-ABL1 kinase domain mutations identified by next-generation sequencing in patients with chronic myeloid leukaemia: a population-based study. Lancet Haematol 2019;6:e276-e284. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/31036317>.

65. Soverini S, Abruzzese E, Bocchia M, et al. Next-generation sequencing for BCR-ABL1 kinase domain mutation testing in patients with chronic myeloid leukemia: a position paper. J Hematol Oncol 2019;12:131. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/31801582>.

66. Soverini S, Bavaro L, De Benedittis C, et al. Prospective assessment of NGS-detectable mutations in CML patients with nonoptimal response: the NEXT-in-CML study. Blood 2020;135:534-541. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/31877211>.

67. Sokal J, Cox E, Baccarani M, et al. Prognostic discrimination in "good-risk" chronic granulocytic leukemia. Blood 1984;63:789-799. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/6584184>.

68. Hasford J, Pffirmann M, Hehlmann R, et al. A new prognostic score for survival of patients with chronic myeloid leukemia treated with interferon alfa. Writing Committee for the Collaborative CML Prognostic Factors Project Group. J Natl Cancer Inst 1998;90:850-858. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/9625174>.

69. Pffirmann M, Baccarani M, Saussele S, et al. Prognosis of long-term survival considering disease-specific death in patients with chronic myeloid leukemia. Leukemia 2016;30:48-56. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/26416462>.

70. Cortes JE, Talpaz M, O'Brien S, et al. Staging of chronic myeloid leukemia in the imatinib era: an evaluation of the World Health Organization proposal. Cancer 2006;106:1306-1315. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/16463391>.

71. Arber DA, Orazi A, Hasserjian R, et al. The 2016 revision to the World Health Organization classification of myeloid neoplasms and acute

leukemia. Blood 2016;127:2391-2405. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/27069254>.

72. O'Dwyer ME, Mauro MJ, Kurilik G, et al. The impact of clonal evolution on response to imatinib mesylate (STI571) in accelerated phase CML. Blood 2002;100:1628-1633. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/12176881>.

73. Druker BJ. Chronic myelogenous leukemia In: DeVita VT, Lawrence TS, Rosenberg SA, eds. DeVita, Hellman, and Rosenberg's Cancer: Principles & Practice of Oncology. Vol. 2 (ed 8): Lippincott, Williams and Wilkins; 2007:2267-2304.

74. Hochhaus A, Larson RA, Guilhot F, et al. Long-term outcomes of imatinib treatment for chronic myeloid leukemia. N Engl J Med 2017;376:917-927. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/28273028>.

75. Cortes JE, Saglio G, Kantarjian HM, et al. Final 5-year study results of DASISION: the dasatinib versus imatinib study in treatment-naïve chronic myeloid leukemia patients trial. J Clin Oncol 2016;34:2333-2340. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/27217448>.

76. Hochhaus A, Saglio G, Hughes TP, et al. Long-term benefits and risks of frontline nilotinib vs imatinib for chronic myeloid leukemia in chronic phase: 5-year update of the randomized ENESTnd trial. Leukemia 2016;30:1044-1054. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/26837842>.

77. Cortes JE, Gambacorti-Passerini C, Deininger MW, et al. Bosutinib Versus Imatinib for Newly Diagnosed Chronic Myeloid Leukemia: Results From the Randomized BFORE Trial. J Clin Oncol 2018;36:231-237. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/29091516>.

78. Baccarani M, Druker BJ, Branford S, et al. Long-term response to imatinib is not affected by the initial dose in patients with Philadelphia chromosome-positive chronic myeloid leukemia in chronic phase: final update from the Tyrosine Kinase Inhibitor Optimization and Selectivity (TOPS) study. Int J Hematol 2014;99:616-624. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/24658916>.



NCCN Guidelines Version 2.2022

Chronic Myeloid Leukemia

79. Deininger MW, Kopecky KJ, Radich JP, et al. Imatinib 800 mg daily induces deeper molecular responses than imatinib 400 mg daily: results of SWOG S0325, an intergroup randomized PHASE II trial in newly diagnosed chronic phase chronic myeloid leukaemia. *Br J Haematol* 2014;164:223-232. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/24383843>.

80. Hehlmann R, Lauseker M, Saussele S, et al. Assessment of imatinib as first-line treatment of chronic myeloid leukemia: 10-year survival results of the randomized CML study IV and impact of non-CML determinants. *Leukemia* 2017;31:2398-2406. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/28804124>.

81. Hoffmann VS, Hasford J, Deininger M, et al. Systematic review and meta-analysis of standard-dose imatinib vs. high-dose imatinib and second generation tyrosine kinase inhibitors for chronic myeloid leukemia. *J Cancer Res Clin Oncol* 2017;143:1311-1318. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/28275866>.

82. Quintas-Cardama A, Han X, Kantarjian H, Cortes J. Tyrosine kinase inhibitor-induced platelet dysfunction in patients with chronic myeloid leukemia. *Blood* 2009;114:261-263. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/19414863>.

83. Hughes TP, Laneuville P, Rousselot P, et al. Incidence, outcomes, and risk factors of pleural effusion in patients receiving dasatinib therapy for Philadelphia chromosome-positive leukemia. *Haematologica* 2019;104:93-101. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/30093398>.

84. Porkka K, Khoury HJ, Paquette RL, et al. Dasatinib 100 mg once daily minimizes the occurrence of pleural effusion in patients with chronic myeloid leukemia in chronic phase and efficacy is unaffected in patients who develop pleural effusion. *Cancer* 2010;116:377-386. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/19924787>.

85. Montani D, Bergot E, Gunther S, et al. Pulmonary arterial hypertension in patients treated by dasatinib. *Circulation* 2012;125:2128-2137. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/22451584>.

86. Orlandi EM, Rocca B, Pazzano AS, Ghio S. Reversible pulmonary arterial hypertension likely related to long-term, low-dose dasatinib

treatment for chronic myeloid leukaemia. *Leuk Res* 2012;36:e4-6.

Available at: <http://www.ncbi.nlm.nih.gov/pubmed/21890201>.

87. Cirmi S, El Abd A, Letinier L, et al. Cardiovascular Toxicity of Tyrosine Kinase Inhibitors Used in Chronic Myeloid Leukemia: An Analysis of the FDA Adverse Event Reporting System Database (FAERS). *Cancers (Basel)* 2020;12. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/32235443>.

88. Naqvi K, Jabbour E, Skinner J, et al. Long-term follow-up of lower dose dasatinib (50 mg daily) as frontline therapy in newly diagnosed chronic-phase chronic myeloid leukemia. *Cancer* 2020;126:67-75. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/31553487>.

89. Bergeron A, Rea D, Levy V, et al. Lung abnormalities after dasatinib treatment for chronic myeloid leukemia: a case series. *Am J Respir Crit Care Med* 2007;176:814-818. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/17600277>.

90. Serpa M, Sanabani SS, Bendit I, et al. Efficacy and tolerability after unusually low doses of dasatinib in chronic myeloid leukemia patients intolerant to standard-dose dasatinib therapy. *Clin Med Insights Oncol* 2010;4:155-162. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/21234296>.

91. Efficace F, Baccarani M, Breccia M, et al. Chronic fatigue is the most important factor limiting health-related quality of life of chronic myeloid leukemia patients treated with imatinib. *Leukemia* 2013;27:1511-1519. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/23417029>.

92. Berman E, Nicolaidis M, Maki RG, et al. Altered bone and mineral metabolism in patients receiving imatinib mesylate. *N Engl J Med* 2006;354:2006-2013. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/16687713>.

93. Berman E, Girotra M, Cheng C, et al. Effect of long term imatinib on bone in adults with chronic myelogenous leukemia and gastrointestinal stromal tumors. *Leuk Res* 2013;37:790-794. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/23473999>.



NCCN Guidelines Version 2.2022

Chronic Myeloid Leukemia

94. Tsao AS, Kantarjian H, Cortes J, et al. Imatinib mesylate causes hypopigmentation in the skin. *Cancer* 2003;98:2483-2487. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/14635084>.
95. Aleem A. Hypopigmentation of the skin due to imatinib mesylate in patients with chronic myeloid leukemia. *Hematol Oncol Stem Cell Ther* 2009;2:358-361. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/20118061>.
96. Sakurai M, Kikuchi T, Karigane D, et al. Renal dysfunction and anemia associated with long-term imatinib treatment in patients with chronic myelogenous leukemia. *Int J Hematol* 2019;109:292-298. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/30680668>.
97. Aichberger KJ, Herndlhofer S, Schernthaner G-H, et al. Progressive peripheral arterial occlusive disease and other vascular events during nilotinib therapy in CML. *Am J Hematol* 2011;86:533-539. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/21538470>.
98. Tefferi A, Letendre L. Nilotinib treatment-associated peripheral artery disease and sudden death: yet another reason to stick to imatinib as front-line therapy for chronic myelogenous leukemia. *Am J Hematol* 2011;86:610-611. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/21630307>.
99. Giles FJ, Mauro MJ, Hong F, et al. Rates of peripheral arterial occlusive disease in patients with chronic myeloid leukemia in the chronic phase treated with imatinib, nilotinib, or non-tyrosine kinase therapy: a retrospective cohort analysis. *Leukemia* 2013;27:1310-1315. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/23459450>.
100. Assuncao PM, Lana TP, Delamain MT, et al. Cardiovascular Risk and Cardiovascular Events in Patients With Chronic Myeloid Leukemia Treated With Tyrosine Kinase Inhibitors. *Clin Lymphoma Myeloma Leuk* 2019;19:162-166. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/30660512>.
101. Caocci G, Mulas O, Bonifacio M, et al. Recurrent arterial occlusive events in patients with chronic myeloid leukemia treated with second- and third-generation tyrosine kinase inhibitors and role of secondary prevention. *Int J Cardiol* 2019;288:124-127. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/31029498>.
102. Tokuhira M, Kimura Y, Sugimoto K, et al. Efficacy and safety of nilotinib therapy in patients with newly diagnosed chronic myeloid leukemia in the chronic phase. *Med Oncol* 2018;35:38. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/29442179>.
103. Quintas-Cardama A, Kantarjian H, O'Brien S, et al. Granulocyte-colony-stimulating factor (filgrastim) may overcome imatinib-induced neutropenia in patients with chronic-phase chronic myelogenous leukemia. *Cancer* 2004;100:2592-2597. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/15197801>.
104. Quintas-Cardama A, De Souza Santos FP, Kantarjian H, et al. Dynamics and management of cytopenias associated with dasatinib therapy in patients with chronic myeloid leukemia in chronic phase after imatinib failure. *Cancer* 2009;115:3935-3943. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/19517473>.
105. Santos FP, Alvarado Y, Kantarjian H, et al. Long-term prognostic impact of the use of erythropoietic-stimulating agents in patients with chronic myeloid leukemia in chronic phase treated with imatinib. *Cancer* 2011;117:982-991. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/20960502>.
106. Reinhold U, Hennig E, Leiblein S, et al. FISH for BCR-ABL on interphases of peripheral blood neutrophils but not of unselected white cells correlates with bone marrow cytogenetics in CML patients treated with imatinib. *Leukemia* 2003;17:1925-1929. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/14513039>.
107. Fugazza G, Miglino M, Bruzzone R, et al. Cytogenetic and fluorescence in situ hybridization monitoring in Ph+ Chronic Myeloid Leukemia patients treated with imatinib mesylate. *J Exp Clin Cancer Res* 2004;23:295-299. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/15354415>.
108. Landstrom AP, Ketterling RP, Knudson RA, Tefferi A. Utility of peripheral blood dual color, double fusion fluorescent in situ hybridization for BCR/ABL fusion to assess cytogenetic remission status in chronic myeloid leukemia. *Leuk Lymphoma* 2006;47:2055-2061. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/17071476>.



NCCN Guidelines Version 2.2022

Chronic Myeloid Leukemia

109. Testoni N, Marzocchi G, Luatti S, et al. Chronic myeloid leukemia: a prospective comparison of interphase fluorescence in situ hybridization and chromosome banding analysis for the definition of complete cytogenetic response: a study of the GIMEMA CML WP. *Blood* 2009;114:4939-4943. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/19797518>.

110. Lima L, Bernal-Mizrachi L, Saxe D, et al. Peripheral blood monitoring of chronic myeloid leukemia during treatment with imatinib, second-line agents, and beyond. *Cancer* 2011;117:1245-1252. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/21381013>.

111. Hughes T, Hochhaus A, Branford S, et al. Long-term prognostic significance of early molecular response to imatinib in newly diagnosed chronic myeloid leukemia: an analysis from the International Randomized Study of Interferon and STI571 (IRIS). *Blood* 2010;116:3758-3765. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/20679528>.

112. Akard LP, Cortes JE, Albitar M, et al. Correlations between cytogenetic and molecular monitoring among patients with newly diagnosed chronic myeloid leukemia in chronic phase: post hoc analyses of the rationale and insight for gleevec high-dose therapy study. *Arch Pathol Lab Med* 2014;138:1186-1192. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/24308645>.

113. Branford S, Cross NCP, Hochhaus A, et al. Rationale for the recommendations for harmonizing current methodology for detecting BCR-ABL transcripts in patients with chronic myeloid leukaemia. *Leukemia* 2006;20:1925-1930. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/16990771>.

114. Cross NC. Standardisation of molecular monitoring for chronic myeloid leukaemia. *Best Pract Res Clin Haematol* 2009;22:355-365. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/19959086>.

115. Cross NC, White HE, Muller MC, et al. Standardized definitions of molecular response in chronic myeloid leukemia. *Leukemia* 2012;26:2172-2175. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/22504141>.

116. Branford S, Fletcher L, Cross NC, et al. Desirable performance characteristics for BCR-ABL measurement on an international reporting scale to allow consistent interpretation of individual patient response and

comparison of response rates between clinical trials. *Blood* 2008;112:3330-3338. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/18684859>.

117. Guerin A, Chen L, Dea K, et al. Association between regular molecular monitoring and tyrosine kinase inhibitor therapy adherence in chronic myelogenous leukemia in the chronic phase. *Curr Med Res Opin* 2014;30:1345-1352. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/24640967>.

118. Hanfstein B, Muller MC, Hehlmann R, et al. Early molecular and cytogenetic response is predictive for long-term progression-free and overall survival in chronic myeloid leukemia (CML). *Leukemia* 2012;26:2096-2102. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/22446502>.

119. Marin D, Ibrahim AR, Lucas C, et al. Assessment of BCR-ABL1 transcript levels at 3 months is the only requirement for predicting outcome for patients with chronic myeloid leukemia treated with tyrosine kinase inhibitors. *J Clin Oncol* 2012;30:232-238. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/22067393>.

120. Neelakantan P, Gerrard G, Lucas C, et al. Combining BCR-ABL1 transcript levels at 3 and 6 months in chronic myeloid leukemia: implications for early intervention strategies. *Blood* 2013;121:2739-2742. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/23380743>.

121. Nazha A, Kantarjian H, Jain P, et al. Assessment at 6 months may be warranted for patients with chronic myeloid leukemia with no major cytogenetic response at 3 months. *Haematologica* 2013;98:1686-1688. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/23812943>.

122. Branford S, Yeung DT, Parker WT, et al. Prognosis for patients with CML and >10% BCR-ABL1 after 3 months of imatinib depends on the rate of BCR-ABL1 decline. *Blood* 2014;124:511-518. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/24859364>.

123. Hanfstein B, Shlyakhto V, Lauseker M, et al. Velocity of early BCR-ABL transcript elimination as an optimized predictor of outcome in chronic myeloid leukemia (CML) patients in chronic phase on treatment with imatinib. *Leukemia* 2014;28:1988-1992. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/24798484>.



NCCN Guidelines Version 2.2022

Chronic Myeloid Leukemia

124. Iriyama N, Fujisawa S, Yoshida C, et al. Shorter halving time of BCR-ABL1 transcripts is a novel predictor for achievement of molecular responses in newly diagnosed chronic-phase chronic myeloid leukemia treated with dasatinib: Results of the D-first study of Kanto CML study group. *Am J Hematol* 2015;90:282-287. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/25530131>.

125. Hochhaus A, O'Brien SG, Guilhot F, et al. Six-year follow-up of patients receiving imatinib for the first-line treatment of chronic myeloid leukemia. *Leukemia* 2009;23:1054-1061. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/19282833>.

126. Jabbour E, Kantarjian H, O'Brien S, et al. The achievement of an early complete cytogenetic response is a major determinant for outcome in patients with early chronic phase chronic myeloid leukemia treated with tyrosine kinase inhibitors. *Blood* 2011;118:4541-4546. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/21803854>.

127. Druker BJ, Guilhot F, O'Brien SG, et al. Five-year follow-up of patients receiving imatinib for chronic myeloid leukemia. *N Engl J Med* 2006;355:2408-2417. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/17151364>.

128. Press RD, Galderisi C, Yang R, et al. A half-log increase in BCR-ABL RNA predicts a higher risk of relapse in patients with chronic myeloid leukemia with an imatinib-induced complete cytogenetic response. *Clin Cancer Res* 2007;13:6136-6143. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/17947479>.

129. de Lavallade H, Apperley JF, Khorashad JS, et al. Imatinib for newly diagnosed patients with chronic myeloid leukemia: incidence of sustained responses in an intention-to-treat analysis. *J Clin Oncol* 2008;26:3358-3363. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/18519952>.

130. Marin D, Milojkovic D, Olavarria E, et al. European LeukemiaNet criteria for failure or suboptimal response reliably identify patients with CML in early chronic phase treated with imatinib whose eventual outcome is poor. *Blood* 2008;112:4437-4444. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/18716134>.

131. Jabbour E, Kantarjian HM, O'Brien S, et al. Front-line therapy with second-generation tyrosine kinase inhibitors in patients with early chronic

phase chronic myeloid leukemia: what is the optimal response? *J Clin Oncol* 2011;29:4260-4265. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/21990394>.

132. Hehlmann R, Muller MC, Lauseker M, et al. Deep molecular response is reached by the majority of patients treated with imatinib, predicts survival, and is achieved more quickly by optimized high-dose imatinib: results from the randomized CML-study IV. *J Clin Oncol* 2014;32:415-423. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/24297946>.

133. Saussele S, Hehlmann R, Fabarius A, et al. Defining therapy goals for major molecular remission in chronic myeloid leukemia: results of the randomized CML Study IV. *Leukemia* 2018;32:1222-1228. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/29479070>.

134. Cervantes F, López-Garrido P, Montero MI, et al. Early intervention during imatinib therapy in patients with newly diagnosed chronic-phase chronic myeloid leukemia: a study of the Spanish PETHEMA group. *Haematologica* 2010;95:1317-1324. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/20220063>.

135. Kantarjian H, Cortes J. Considerations in the management of patients with Philadelphia chromosome-positive chronic myeloid leukemia receiving tyrosine kinase inhibitor therapy. *J Clin Oncol* 2011;29:1512-1516. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/21422414>.

136. Hochhaus A, Baccarani M, Silver RT, et al. European LeukemiaNet 2020 recommendations for treating chronic myeloid leukemia. *Leukemia* 2020;34:966-984. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/32127639>.

137. Shah NP, Rousselot P, Schiffer C, et al. Dasatinib in imatinib-resistant or -intolerant chronic-phase, chronic myeloid leukemia patients: 7-year follow-up of study CA180-034. *Am J Hematol* 2016;91:869-874. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/27192969>.

138. Giles FJ, le Coutre PD, Pinilla-Ibarz J, et al. Nilotinib in imatinib-resistant or imatinib-intolerant patients with chronic myeloid leukemia in chronic phase: 48-month follow-up results of a phase II study. *Leukemia* 2013;27:107-112. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/22763385>.



NCCN Guidelines Version 2.2022

Chronic Myeloid Leukemia

139. Gambacorti-Passerini C, Cortes JE, Lipton JH, et al. Safety and efficacy of second-line bosutinib for chronic phase chronic myeloid leukemia over a five-year period: final results of a phase I/II study. *Haematologica* 2018;103:1298-1307. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/29773593>.

140. Cortes JE, Kim D-W, Pinilla-Ibarz J, et al. Ponatinib efficacy and safety in Philadelphia chromosome–positive leukemia: final 5-year results of the phase 2 PACE trial. *Blood* 2018;132:393-404. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/29567798>.

141. Kantarjian HM, Talpaz M, O'Brien S, et al. Dose escalation of imatinib mesylate can overcome resistance to standard-dose therapy in patients with chronic myelogenous leukemia. *Blood* 2003;101:473-475. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/12393385>.

142. Marin D, Goldman JM, Olavarria E, Apperley JF. Transient benefit only from increasing the imatinib dose in CML patients who do not achieve complete cytogenetic remissions on conventional doses. *Blood* 2003;102:2702-2704. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/14504074>.

143. Jabbour E, Kantarjian HM, Jones D, et al. Imatinib mesylate dose escalation is associated with durable responses in patients with chronic myeloid leukemia after cytogenetic failure on standard-dose imatinib therapy. *Blood* 2009;113:2154-2160. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/19060245>.

144. Kantarjian HM, Larson RA, Guilhot F, et al. Efficacy of imatinib dose escalation in patients with chronic myeloid leukemia in chronic phase. *Cancer* 2009;115:551-560. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/19117345>.

145. Yeung DT, Osborn MP, White DL, et al. TIDEL-II: first-line use of imatinib in CML with early switch to nilotinib for failure to achieve time-dependent molecular targets. *Blood* 2015;125:915-923. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/25519749>.

146. Cortes JE, De Souza CA, Ayala M, et al. Switching to nilotinib versus imatinib dose escalation in patients with chronic myeloid leukaemia in chronic phase with suboptimal response to imatinib (LASOR): a

randomised, open-label trial. *Lancet Haematol* 2016;3:e581-e591. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/27890073>.

147. Cortes JE, Khoury HJ, Kantarjian HM, et al. Long-term bosutinib for chronic phase chronic myeloid leukemia after failure of imatinib plus dasatinib and/or nilotinib. *Am J Hematol* 2016;91:1206-1214. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/27531525>.

148. Garcia-Gutierrez V, Milojkovic D, Hernandez-Boluda JC, et al. Safety and efficacy of bosutinib in fourth-line therapy of chronic myeloid leukemia patients. *Ann Hematol* 2019;98:321-330. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/30446802>.

149. Jain P, Kantarjian H, Boddur PC, et al. Analysis of cardiovascular and arteriothrombotic adverse events in chronic-phase CML patients after frontline TKIs. *Blood Adv* 2019;3:851-861. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/30885996>.

150. Caocci G, Mulas O, Abruzzese E, et al. Arterial occlusive events in chronic myeloid leukemia patients treated with ponatinib in the real-life practice are predicted by the Systematic Coronary Risk Evaluation (SCORE) chart. *Hematol Oncol* 2019;37:296-302. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/30892724>.

151. Cortes JE, Kantarjian H, Shah NP, et al. Ponatinib in refractory Philadelphia chromosome–positive leukemias. *N Engl J Med* 2012;367:2075-2088. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/23190221>.

152. Dorer DJ, Knickerbocker RK, Baccarani M, et al. Impact of dose intensity of ponatinib on selected adverse events: Multivariate analyses from a pooled population of clinical trial patients. *Leuk Res* 2016;48:84-91. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/27505637>.

153. Iurlo A, Cattaneo D, Orofino N, et al. Low-Dose Ponatinib in Intolerant Chronic Myeloid Leukemia Patients: A Safe and Effective Option. *Clin Drug Investig* 2018;38:475-476. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/29363026>.

154. Casavecchia G, Galderisi M, Novo G, et al. Early diagnosis, clinical management, and follow-up of cardiovascular events with ponatinib. *Heart*



NCCN Guidelines Version 2.2022

Chronic Myeloid Leukemia

Fail Rev 2020. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/32026180>.

155. Garg RJ, Kantarjian H, O'Brien S, et al. The use of nilotinib or dasatinib after failure to 2 prior tyrosine kinase inhibitors: long-term follow-up. *Blood* 2009;114:4361-4368. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/19729517>.

156. Cortes J, Lipton JH, Rea D, et al. Phase 2 study of subcutaneous omacetaxine mepesuccinate after TKI failure in patients with chronic-phase CML with T315I mutation. *Blood* 2012;120:2573-2580. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/22896000>.

157. Cortes J, Digumarti R, Parikh PM, et al. Phase 2 study of subcutaneous omacetaxine mepesuccinate for chronic-phase chronic myeloid leukemia patients resistant to or intolerant of tyrosine kinase inhibitors. *Am J Hematol* 2013;88:350-354. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/23468307>.

158. van Leeuwen RW, van Gelder T, Mathijssen RH, Jansman FG. Drug-drug interactions with tyrosine-kinase inhibitors: a clinical perspective. *Lancet Oncol* 2014;15:e315-326. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/24988935>.

159. Osorio S, Escudero-Vilaplana V, Gomez-Centurion I, et al. Drug-to-drug interactions of tyrosine kinase inhibitors in chronic myeloid leukemia patients. Is it a real problem? *Ann Hematol* 2018;97:2089-2098. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/29955943>.

160. Noens L, van Lierde M-A, De Bock R, et al. Prevalence, determinants, and outcomes of nonadherence to imatinib therapy in patients with chronic myeloid leukemia: the ADAGIO study. *Blood* 2009;113:5401-5411. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/19349618>.

161. Marin D, Bazeos A, Mahon F-X, et al. Adherence is the critical factor for achieving molecular responses in patients with chronic myeloid leukemia who achieve complete cytogenetic responses on imatinib. *J Clin Oncol* 2010;28:2381-2388. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/20385986>.

162. Ibrahim AR, Eliasson L, Apperley JF, et al. Poor adherence is the main reason for loss of CCyR and imatinib failure for chronic myeloid leukemia patients on long-term therapy. *Blood* 2011;117:3733-3736. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/21346253>.

163. Wu EQ, Guerin A, Yu AP, et al. Retrospective real-world comparison of medical visits, costs, and adherence between nilotinib and dasatinib in chronic myeloid leukemia. *Curr Med Res Opin* 2010;26:2861-2869. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/21062136>.

164. Yood MU, Oliveria SA, Cziraky M, et al. Adherence to treatment with second-line therapies, dasatinib and nilotinib, in patients with chronic myeloid leukemia. *Curr Med Res Opin* 2012;28:213-219. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/22168217>.

165. Quintas-Cardama A, Cortes JE, Kantarjian H. Practical management of toxicities associated with tyrosine kinase inhibitors in chronic myeloid leukemia. *Clin Lymphoma Myeloma* 2008;8 Suppl 3:S82-88. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/19254885>.

166. Cornelison M, Jabbour EJ, Welch MA. Managing side effects of tyrosine kinase inhibitor therapy to optimize adherence in patients with chronic myeloid leukemia: the role of the midlevel practitioner. *J Support Oncol* 2012;10:14-24. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/22244674>.

167. Cortes JE, Lipton JH, Miller CB, et al. Evaluating the Impact of a Switch to Nilotinib on Imatinib-Related Chronic Low-Grade Adverse Events in Patients With CML-CP: The ENRICH Study. *Clin Lymphoma Myeloma Leuk* 2016;16:286-296. Available at:

168. Kim DW, Saussele S, Williams LA, et al. Outcomes of switching to dasatinib after imatinib-related low-grade adverse events in patients with chronic myeloid leukemia in chronic phase: the DASPERSE study. *Ann Hematol* 2018;97:1357-1367. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/29556695>.

169. Hiwase D, Tan P, D'Rozario J, et al. Efficacy and safety of nilotinib 300mg twice daily in patients with chronic myeloid leukemia in chronic phase who are intolerant to prior tyrosine kinase inhibitors: Results from the Phase IIIb ENESTswift study. *Leuk Res* 2018;67:109-115. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/29494928>.



NCCN Guidelines Version 2.2022

Chronic Myeloid Leukemia

170. Thomas J, Wang L, Clark RE, Pirmohamed M. Active transport of imatinib into and out of cells: implications for drug resistance. *Blood* 2004;104:3739-3745. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/15315971>.

171. Mahon FX, Hayette S, Lagarde V, et al. Evidence that resistance to nilotinib may be due to BCR-ABL, Pgp, or Src kinase overexpression. *Cancer Res* 2008;68:9809-9816. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/19047160>.

172. Hegedus C, Ozvegy-Laczka C, Apati A, et al. Interaction of nilotinib, dasatinib and bosutinib with ABCB1 and ABCG2: implications for altered anti-cancer effects and pharmacological properties. *Br J Pharmacol* 2009;158:1153-1164. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/19785662>.

173. Picard S, Titier K, Etienne G, et al. Trough imatinib plasma levels are associated with both cytogenetic and molecular responses to standard-dose imatinib in chronic myeloid leukemia. *Blood* 2007;109:3496-3499. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/17192396>.

174. Larson RA, Druker BJ, Guilhot F, et al. Imatinib pharmacokinetics and its correlation with response and safety in chronic-phase chronic myeloid leukemia: a subanalysis of the IRIS study. *Blood* 2008;111:4022-4028. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/18256322>.

175. Bouchet S, Titier K, Moore N, et al. Therapeutic drug monitoring of imatinib in chronic myeloid leukemia: experience from 1216 patients at a centralized laboratory. *Fundam Clin Pharmacol* 2013;27:690-697. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/23113675>.

176. White DL, Radich J, Soverini S, et al. Chronic phase chronic myeloid leukemia patients with low OCT-1 activity randomised to high-dose imatinib achieve better responses, and lower failure rates, than those randomized to standard-dose. *Haematologica* 2012;97:907-914. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/22207690>.

177. Giannoudis A, Davies A, Lucas CM, et al. Effective dasatinib uptake may occur without human organic cation transporter 1 (hOCT1): implications for the treatment of imatinib-resistant chronic myeloid leukemia. *Blood* 2008;112:3348-3354. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/18669873>.

178. Hiwase DK, Saunders V, Hewett D, et al. Dasatinib cellular uptake and efflux in chronic myeloid leukemia cells: therapeutic implications. *Clin Cancer Res* 2008;14:3881-3888. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/18559609>.

179. Davies A, Jordanides NE, Giannoudis A, et al. Nilotinib concentration in cell lines and primary CD34(+) chronic myeloid leukemia cells is not mediated by active uptake or efflux by major drug transporters. *Leukemia* 2009;23:1999-2006. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/19710702>.

180. White DL, Saunders VA, Dang P, et al. OCT-1-mediated influx is a key determinant of the intracellular uptake of imatinib but not nilotinib (AMN107): reduced OCT-1 activity is the cause of low in vitro sensitivity to imatinib. *Blood* 2006;108:697-704. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/16597591>.

181. Branford S, Rudzki Z, Walsh S, et al. Detection of BCR-ABL mutations in patients with CML treated with imatinib is virtually always accompanied by clinical resistance, and mutations in the ATP phosphate-binding loop (P-loop) are associated with a poor prognosis. *Blood* 2003;102:276-283. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/12623848>.

182. Soverini S, Martinelli G, Rosti G, et al. ABL mutations in late chronic phase chronic myeloid leukemia patients with up-front cytogenetic resistance to imatinib are associated with a greater likelihood of progression to blast crisis and shorter survival: a study by the GIMEMA Working Party on Chronic Myeloid Leukemia. *J Clin Oncol* 2005;23:4100-4109. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/15867198>.

183. Nicolini FE, Corm S, Le QH, et al. Mutation status and clinical outcome of 89 imatinib mesylate-resistant chronic myelogenous leukemia patients: a retrospective analysis from the French intergroup of CML (Fi(phi)-LMC GROUP). *Leukemia* 2006;20:1061-1106. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/16642048>.

184. Soverini S, Colarossi S, Gnani A, et al. Contribution of ABL kinase domain mutations to imatinib resistance in different subsets of Philadelphia-positive patients: by the GIMEMA Working Party on Chronic



NCCN Guidelines Version 2.2022

Chronic Myeloid Leukemia

Myeloid Leukemia. Clin Cancer Res 2006;12:7374-7379. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/17189410>.

185. Khorashad JS, de Lavallade H, Apperley JF, et al. Finding of kinase domain mutations in patients with chronic phase chronic myeloid leukemia responding to imatinib may identify those at high risk of disease progression. J Clin Oncol 2008;26:4806-4813. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/18645191>.

186. Soverini S, Gnani A, Colarossi S, et al. Philadelphia-positive patients who already harbor imatinib-resistant Bcr-Abl kinase domain mutations have a higher likelihood of developing additional mutations associated with resistance to second- or third-line tyrosine kinase inhibitors. Blood 2009;114:2168-2171. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/19589924>.

187. Nicolini FE, Hayette S, Corm S, et al. Clinical outcome of 27 imatinib mesylate-resistant chronic myelogenous leukemia patients harboring a T315I BCR-ABL mutation. Haematologica 2007;92:1238-1241. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/17768119>.

188. Jabbour E, Kantarjian H, Jones D, et al. Characteristics and outcomes of patients with chronic myeloid leukemia and T315I mutation following failure of imatinib mesylate therapy. Blood 2008;112:53-55. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/18403620>.

189. Soverini S, Colarossi S, Gnani A, et al. Resistance to dasatinib in Philadelphia-positive leukemia patients and the presence or the selection of mutations at residues 315 and 317 in the BCR-ABL kinase domain. Haematologica 2007;92:401-404. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/17339191>.

190. Jabbour E, Kantarjian HM, Jones D, et al. Characteristics and outcome of chronic myeloid leukemia patients with F317L BCR-ABL kinase domain mutation after therapy with tyrosine kinase inhibitors. Blood 2008;112:4839-4842. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/18818391>.

191. Muller MC, Cortes JE, Kim D-W, et al. Dasatinib treatment of chronic-phase chronic myeloid leukemia: analysis of responses according to preexisting BCR-ABL mutations. Blood 2009;114:4944-4953. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/19779040>.

192. Hughes T, Saglio G, Branford S, et al. Impact of baseline BCR-ABL mutations on response to nilotinib in patients with chronic myeloid leukemia in chronic phase. J Clin Oncol 2009;27:4204-4210. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/19652056>.

193. Naqvi K, Cortes JE, Luthra R, et al. Characteristics and outcome of chronic myeloid leukemia patients with E255K/V BCR-ABL kinase domain mutations. Int J Hematol 2018;107:689-695. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/29464484>.

194. Khoury HJ, Cortes JE, Kantarjian HM, et al. Bosutinib is active in chronic phase chronic myeloid leukemia after imatinib and dasatinib and/or nilotinib therapy failure. Blood 2012;119:3403-3412. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/22371878>.

195. Khorashad JS, Kelley TW, Szankasi P, et al. BCR-ABL1 compound mutations in tyrosine kinase inhibitor-resistant CML: frequency and clonal relationships. Blood 2013;121:489-498. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/23223358>.

196. Zabriskie MS, Eide CA, Tantravahi SK, et al. BCR-ABL1 compound mutations combining key kinase domain positions confer clinical resistance to ponatinib in Ph chromosome-positive leukemia. Cancer Cell 2014;26:428-442. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/25132497>.

197. Deininger MW, Hodgson JG, Shah NP, et al. Compound mutations in BCR-ABL1 are not major drivers of primary or secondary resistance to ponatinib in CP-CML patients. Blood 2016;127:703-712. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/26603839>.

198. Soverini S, Branford S, Nicolini FE, et al. Implications of BCR-ABL1 kinase domain-mediated resistance in chronic myeloid leukemia. Leuk Res 2014;38:10-20. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/24131888>.

199. Grossmann V, Kohlmann A, Zenger M, et al. A deep-sequencing study of chronic myeloid leukemia patients in blast crisis (BC-CML) detects mutations in 76.9% of cases. Leukemia 2011;25:557-560. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/21274004>.



NCCN Guidelines Version 2.2022

Chronic Myeloid Leukemia

200. Schmidt M, Rinke J, Schafer V, et al. Molecular-defined clonal evolution in patients with chronic myeloid leukemia independent of the BCR-ABL status. *Leukemia* 2014;28:2292-2299. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/25212276>.
201. Soverini S, de Benedittis C, Mancini M, Martinelli G. Mutations in the BCR-ABL1 Kinase Domain and Elsewhere in Chronic Myeloid Leukemia. *Clin Lymphoma Myeloma Leuk* 2015;15 Suppl:S120-128. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/26297264>.
202. Kim T, Tyndel MS, Kim HJ, et al. Spectrum of somatic mutation dynamics in chronic myeloid leukemia following tyrosine kinase inhibitor therapy. *Blood* 2017;129:38-47. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/27733357>.
203. Togasaki E, Takeda J, Yoshida K, et al. Frequent somatic mutations in epigenetic regulators in newly diagnosed chronic myeloid leukemia. *Blood Cancer J* 2017;7:e559. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/28452984>.
204. Branford S, Wang P, Yeung DT, et al. Integrative genomic analysis reveals cancer-associated mutations at diagnosis of CML in patients with high-risk disease. *Blood* 2018;132:948-961. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/29967129>.
205. Branford S, Kim DDH, Apperley JF, et al. Laying the foundation for genomically-based risk assessment in chronic myeloid leukemia. *Leukemia* 2019;33:1835-1850. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/31209280>.
206. Adnan Awad S, Kankainen M, Ojala T, et al. Mutation accumulation in cancer genes relates to nonoptimal outcome in chronic myeloid leukemia. *Blood Adv* 2020;4:546-559. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/32045476>.
207. Branford S, Rudzki Z, Parkinson I, et al. Real-time quantitative PCR analysis can be used as a primary screen to identify patients with CML treated with imatinib who have BCR-ABL kinase domain mutations. *Blood* 2004;104:2926-2932. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/15256429>.
208. Wang L, Knight K, Lucas C, Clark R. The role of serial BCR-ABL transcript monitoring in predicting the emergence of BCR-ABL kinase mutations in imatinib-treated patients with chronic myeloid leukemia. *Haematologica* 2006;91:235-239. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/16461309>.
209. Kantarjian HM, Shan J, Jones D, et al. Significance of increasing levels of minimal residual disease in patients with Philadelphia chromosome-positive chronic myelogenous leukemia in complete cytogenetic response. *J Clin Oncol* 2009;27:3659-3663. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/19487383>.
210. Marin D, Khorashad JS, Foroni L, et al. Does a rise in the BCR-ABL1 transcript level identify chronic phase CML patients responding to imatinib who have a high risk of cytogenetic relapse? *Br J Haematol* 2009;145:373-375. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/19344397>.
211. Press RD, Willis SG, Laudadio J, et al. Determining the rise in BCR-ABL RNA that optimally predicts a kinase domain mutation in patients with chronic myeloid leukemia on imatinib. *Blood* 2009;114:2598-2605. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/19625707>.
212. Mahon FX, Rea D, Guilhot J, et al. Discontinuation of imatinib in patients with chronic myeloid leukaemia who have maintained complete molecular remission for at least 2 years: the prospective, multicentre Stop Imatinib (STIM) trial. *Lancet Oncol* 2010;10:1029-1035. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/20965785>.
213. Etienne G, Guilhot J, Rea D, et al. Long-term follow-up of the french stop imatinib (STIM1) study in patients with chronic myeloid leukemia. *J Clin Oncol* 2017;35:298-305. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/28095277>.
214. Thielen N, van der Holt B, Cornelissen JJ, et al. Imatinib discontinuation in chronic phase myeloid leukaemia patients in sustained complete molecular response: a randomised trial of the Dutch-Belgian Cooperative Trial for Haemato-Oncology (HOVON). *Eur J Cancer* 2013;49:3242-3246. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/23876833>.
215. Rousselot P, Charbonnier A, Cony-Makhoul P, et al. Loss of major molecular response as a trigger for restarting tyrosine kinase inhibitor



NCCN Guidelines Version 2.2022

Chronic Myeloid Leukemia

therapy in patients with chronic-phase chronic myelogenous leukemia who have stopped imatinib after durable undetectable disease. *J Clin Oncol* 2014;32:424-430. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/24323036>.

216. Mori S, Vagge E, le Coutre P, et al. Age and dPCR can predict relapse in CML patients who discontinued imatinib: the ISAV study. *Am J Hematol* 2015;90:910-914. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/26178642>.

217. Lee SE, Choi SY, Song HY, et al. Imatinib withdrawal syndrome and longer duration of imatinib have a close association with a lower molecular relapse after treatment discontinuation: the KID study. *Haematologica* 2016;101:717-723. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/26888022>.

218. Ross DM, Pagani IS, Shanmuganathan N, et al. Long-term treatment-free remission of chronic myeloid leukemia with falling levels of residual leukemic cells. *Leukemia* 2018;32:2572-2579. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/30315232>.

219. Rea D, Nicolini FE, Tulliez M, et al. Discontinuation of dasatinib or nilotinib in chronic myeloid leukemia: interim analysis of the STOP 2G-TKI study. *Blood* 2017;129:846-854. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/27932374>.

220. Ross DM, Masszi T, Gomez Casares MT, et al. Durable treatment-free remission in patients with chronic myeloid leukemia in chronic phase following frontline nilotinib: 96-week update of the ENESTfreedom study. *J Cancer Res Clin Oncol* 2018;144:945-954. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/29468438>.

221. Mahon FX, Boquimpani C, Kim DW, et al. Treatment-Free Remission After Second-Line Nilotinib Treatment in Patients With Chronic Myeloid Leukemia in Chronic Phase: Results From a Single-Group, Phase 2, Open-Label Study. *Ann Intern Med* 2018;168:461-470. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/29459949>.

222. Okada M, Imagawa J, Tanaka H, et al. Final 3-year Results of the Dasatinib Discontinuation Trial in Patients With Chronic Myeloid Leukemia Who Received Dasatinib as a Second-line Treatment. *Clin Lymphoma*

Myeloma Leuk 2018;18:353-360 e351. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/29610029>.

223. Saussele S, Richter J, Guilhot J, et al. Discontinuation of tyrosine kinase inhibitor therapy in chronic myeloid leukaemia (EURO-SKI): a prespecified interim analysis of a prospective, multicentre, non-randomised, trial. *Lancet Oncol* 2018;19:747-757. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/29735299>.

224. Shah NP, Garcia-Gutierrez V, Jimenez-Velasco A, et al. Dasatinib discontinuation in patients with chronic-phase chronic myeloid leukemia and stable deep molecular response: the DASFREE study. *Leuk Lymphoma* 2020;61:650-659. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/31647335>.

225. Kimura S, Imagawa J, Murai K, et al. Treatment-free remission after first-line dasatinib discontinuation in patients with chronic myeloid leukaemia (first-line DADI trial): a single-arm, multicentre, phase 2 trial. *Lancet Haematol* 2020;7:e218-e225. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/31978329>.

226. Legros L, Nicolini FE, Etienne G, et al. Second tyrosine kinase inhibitor discontinuation attempt in patients with chronic myeloid leukemia. *Cancer* 2017;123:4403-4410. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/28743166>.

227. Ilander M, Olsson-Stromberg U, Schlums H, et al. Increased proportion of mature NK cells is associated with successful imatinib discontinuation in chronic myeloid leukemia. *Leukemia* 2017;31:1108-1116. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/27890936>.

228. Talpaz M, Silver RT, Druker BJ, et al. Imatinib induces durable hematologic and cytogenetic responses in patients with accelerated phase chronic myeloid leukemia: results of a phase 2 study. *Blood* 2002;99:1928-1937. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/11877262>.

229. Kantarjian HM, Cortes J, O'Brien S, et al. Imatinib mesylate (STI571) therapy for Philadelphia chromosome-positive chronic myelogenous leukemia in blast phase. *Blood* 2002;99:3547-3553. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/11986206>.



NCCN Guidelines Version 2.2022

Chronic Myeloid Leukemia

230. Sawyers CL, Hochhaus A, Feldman E, et al. Imatinib induces hematologic and cytogenetic responses in patients with chronic myelogenous leukemia in myeloid blast crisis: results of a phase II study. *Blood* 2002;99:3530-3539. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/11986204>.

231. Palandri F, Castagnetti F, Testoni N, et al. Chronic myeloid leukemia in blast crisis treated with imatinib 600 mg: outcome of the patients alive after a 6-year follow-up. *Haematologica* 2008;93:1792-1796. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/18838477>.

232. Palandri F, Castagnetti F, Alimena G, et al. The long-term durability of cytogenetic responses in patients with accelerated phase chronic myeloid leukemia treated with imatinib 600 mg: the GIMEMA CML Working Party experience after a 7-year follow-up. *Haematologica* 2009;94:205-212. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/19144656>.

233. Kantarjian H, Cortes J, Kim DW, et al. Phase 3 study of dasatinib 140 mg once daily versus 70 mg twice daily in patients with chronic myeloid leukemia in accelerated phase resistant or intolerant to imatinib: 15-month median follow-up. *Blood* 2009;113:6322-6329. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/19369231>.

234. Ottmann O, Saglio G, Apperley JF, et al. Long-term efficacy and safety of dasatinib in patients with chronic myeloid leukemia in accelerated phase who are resistant to or intolerant of imatinib. *Blood Cancer J* 2018;8:88. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/30190469>.

235. Saglio G, Hochhaus A, Goh YT, et al. Dasatinib in imatinib-resistant or imatinib-intolerant chronic myeloid leukemia in blast phase after 2 years of follow-up in a phase 3 study: efficacy and tolerability of 140 milligrams once daily and 70 milligrams twice daily. *Cancer* 2010;116:3852-3861. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/20564086>.

236. le Coutre PD, Giles FJ, Hochhaus A, et al. Nilotinib in patients with Ph+ chronic myeloid leukemia in accelerated phase following imatinib resistance or intolerance: 24-month follow-up results. *Leukemia* 2012;26:1189-1194. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/22076466>.

237. Giles FJ, Kantarjian HM, le Coutre PD, et al. Nilotinib is effective in imatinib-resistant or -intolerant patients with chronic myeloid leukemia in blastic phase. *Leukemia* 2012;26:959-962. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/22157807>.

238. Gambacorti-Passerini C, Kantarjian HM, Kim DW, et al. Long-term efficacy and safety of bosutinib in patients with advanced leukemia following resistance/intolerance to imatinib and other tyrosine kinase inhibitors. *Am J Hematol* 2015;90:755-768. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/26040495>.

239. Oki Y, Kantarjian HM, Gharibyan V, et al. Phase II study of low-dose decitabine in combination with imatinib mesylate in patients with accelerated or myeloid blastic phase of chronic myelogenous leukemia. *Cancer* 2007;109:899-906. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/17236224>.

240. Quintas-Cardama A, Kantarjian H, Garcia-Manero G, et al. A pilot study of imatinib, low-dose cytarabine and idarubicin for patients with chronic myeloid leukemia in myeloid blast phase. *Leuk Lymphoma* 2007;48:283-289. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/17325887>.

241. Fruehauf S, Topaly J, Buss EC, et al. Imatinib combined with mitoxantrone/etoposide and cytarabine is an effective induction therapy for patients with chronic myeloid leukemia in myeloid blast crisis. *Cancer* 2007;109:1543-1549. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/17340589>.

242. Deau B, Nicolini FE, Guilhot J, et al. The addition of daunorubicin to imatinib mesylate in combination with cytarabine improves the response rate and the survival of patients with myeloid blast crisis chronic myelogenous leukemia (AFR01 study). *Leuk Res* 2011;35:777-782. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/21145590>.

243. Benjamini O, Dumlao TL, Kantarjian H, et al. Phase II trial of hyper CVAD and dasatinib in patients with relapsed Philadelphia chromosome positive acute lymphoblastic leukemia or blast phase chronic myeloid leukemia. *Am J Hematol* 2014;89:282-287. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/24779033>.



NCCN Guidelines Version 2.2022

Chronic Myeloid Leukemia

244. Strati P, Kantarjian H, Thomas D, et al. HCVAD plus imatinib or dasatinib in lymphoid blastic phase chronic myeloid leukemia. *Cancer* 2014;120:373-380. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/24151050>.

245. Fava C, Kantarjian HM, Jabbour E, et al. Failure to achieve a complete hematologic response at the time of a major cytogenetic response with second-generation tyrosine kinase inhibitors is associated with a poor prognosis among patients with chronic myeloid leukemia in accelerated or blast phase. *Blood* 2009;113:5058-5063. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/19282457>.

246. Khoury HJ, Cortes J, Baccarani M, et al. Omacetaxine mepesuccinate in patients with advanced chronic myeloid leukemia with resistance or intolerance to tyrosine kinase inhibitors. *Leuk Lymphoma* 2015;56:120-127. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/24650054>.

247. Rea D, Etienne G, Nicolini F, et al. First-line imatinib mesylate in patients with newly diagnosed accelerated phase-chronic myeloid leukemia. *Leukemia* 2012;26:2254-2259. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/22460758>.

248. Ohanian M, Kantarjian HM, Quintas-Cardama A, et al. Tyrosine kinase inhibitors as initial therapy for patients with chronic myeloid leukemia in accelerated phase. *Clin Lymphoma Myeloma Leuk* 2014;14:155-162 e151. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/24332214>.

249. Jain P, Kantarjian HM, Ghorab A, et al. Prognostic factors and survival outcomes in patients with chronic myeloid leukemia in blast phase in the tyrosine kinase inhibitor era: Cohort study of 477 patients. *Cancer* 2017;123:4391-4402. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/28743165>.

250. Rajappa S, Uppin SG, Raghunadharao D, et al. Isolated central nervous system blast crisis in chronic myeloid leukemia. *Hematol Oncol* 2004;22:179-181. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/15995975>.

251. Kim HJ, Jung CW, Kim K, et al. Isolated blast crisis in CNS in a patient with chronic myelogenous leukemia maintaining major cytogenetic

response after imatinib. *J Clin Oncol* 2006;24:4028-4029. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/16921058>.

252. Altintas A, Cil T, Kilinc I, et al. Central nervous system blastic crisis in chronic myeloid leukemia on imatinib mesylate therapy: a case report. *J Neurooncol* 2007;84:103-105. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/17318411>.

253. Aftimos P, Nasr F. Isolated CNS lymphoid blast crisis in a patient with imatinib-resistant chronic myelogenous leukemia: case report and review of the literature. *Leuk Res* 2009;33:e178-180. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/19446330>.

254. Porkka K, Koskenvesa P, Lundan T, et al. Dasatinib crosses the blood-brain barrier and is an efficient therapy for central nervous system Philadelphia chromosome-positive leukemia. *Blood* 2008;112:1005-1012. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/18477770>.

255. Hansen JA, Gooley TA, Martin PJ, et al. Bone marrow transplants from unrelated donors for patients with chronic myeloid leukemia. *N Engl J Med* 1998;338:962-968. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/9521984>.

256. Davies SM, DeFor TE, McGlave PB, et al. Equivalent outcomes in patients with chronic myelogenous leukemia after early transplantation of phenotypically matched bone marrow from related or unrelated donors. *Am J Med* 2001;110:339-346. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/11286947>.

257. Crawley C, Szydlo R, Lalancette M, et al. Outcomes of reduced-intensity transplantation for chronic myeloid leukemia: an analysis of prognostic factors from the Chronic Leukemia Working Party of the EBMT. *Blood* 2005;106:2969-2976. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/15998838>.

258. Or R, Shapira MY, Resnick I, et al. Nonmyeloablative allogeneic stem cell transplantation for the treatment of chronic myeloid leukemia in first chronic phase. *Blood* 2003;101:441-445. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/12393604>.

259. Faber E, Koza V, Vitek A, et al. Reduced-intensity conditioning for allogeneic stem cell transplantation in patients with chronic myeloid



NCCN Guidelines Version 2.2022

Chronic Myeloid Leukemia

leukemia is associated with better overall survival but inferior disease-free survival when compared with myeloablative conditioning - a retrospective study of the Czech National Hematopoietic Stem Cell Transplantation Registry. *Neoplasma* 2007;54:443-446. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/17688375>.

260. Kebriaei P, Detry MA, Giralt S, et al. Long-term follow-up of allogeneic hematopoietic stem-cell transplantation with reduced-intensity conditioning for patients with chronic myeloid leukemia. *Blood* 2007;110:3456-3462. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/17652620>.

261. Warlick E, Ahn KW, Pedersen TL, et al. Reduced intensity conditioning is superior to nonmyeloablative conditioning for older chronic myelogenous leukemia patients undergoing hematopoietic cell transplant during the tyrosine kinase inhibitor era. *Blood* 2012;119:4083-4090. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/22408257>.

262. Velev N, Cortes J, Champlin R, et al. Stem cell transplantation for patients with chronic myeloid leukemia resistant to tyrosine kinase inhibitors with BCR-ABL kinase domain mutation T315I. *Cancer* 2010;116:3631-3637. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/20564073>.

263. Jabbour E, Cortes J, Santos FP, et al. Results of allogeneic hematopoietic stem cell transplantation for chronic myelogenous leukemia patients who failed tyrosine kinase inhibitors after developing BCR-ABL1 kinase domain mutations. *Blood* 2011;117:3641-3647. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/21156844>.

264. Nicolini FE, Basak GW, Soverini S, et al. Allogeneic stem cell transplantation for patients harboring T315I BCR-ABL mutated leukemias. *Blood* 2011;118:5697-5700. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/21926354>.

265. Nair AP, Barnett MJ, Broady RC, et al. Allogeneic hematopoietic stem cell transplantation is an effective salvage therapy for patients with chronic myeloid leukemia presenting with advanced disease or failing treatment with tyrosine kinase inhibitors. *Biol Blood Marrow Transplant* 2015;21:1437-1444. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/25865648>.

266. Deininger M, Schleuning M, Greinix H, et al. The effect of prior exposure to imatinib on transplant-related mortality. *Haematologica* 2006;91:452-459. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/16585011>.

267. Oehler VG, Gooley T, Snyder DS, et al. The effects of imatinib mesylate treatment before allogeneic transplantation for chronic myeloid leukemia. *Blood* 2007;109:1782-1789. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/17062727>.

268. Jabbour E, Cortes J, Kantarjian H, et al. Novel tyrosine kinase inhibitor therapy before allogeneic stem cell transplantation in patients with chronic myeloid leukemia: no evidence for increased transplant-related toxicity. *Cancer* 2007;110:340-344. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/17559140>.

269. Shimoni A, Leiba M, Schleuning M, et al. Prior treatment with the tyrosine kinase inhibitors dasatinib and nilotinib allows stem cell transplantation (SCT) in a less advanced disease phase and does not increase SCT Toxicity in patients with chronic myelogenous leukemia and philadelphia positive acute lymphoblastic leukemia. *Leukemia* 2009;23:190-194. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/18596746>.

270. Breccia M, Palandri F, Iori AP, et al. Second-generation tyrosine kinase inhibitors before allogeneic stem cell transplantation in patients with chronic myeloid leukemia resistant to imatinib. *Leuk Res* 2010;34:143-147. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/19481800>.

271. Lee SE, Choi SY, Kim SH, et al. Prognostic factors for outcomes of allogeneic stem cell transplantation in chronic phase chronic myeloid leukemia in the era of tyrosine kinase inhibitors. *Hematology* 2014;19:63-72. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/23684143>.

272. Piekarska A, Gil L, Prejzner W, et al. Pretransplantation use of the second-generation tyrosine kinase inhibitors has no negative impact on the HCT outcome. *Ann Hematol* 2015;94:1891-1897. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/26220759>.

273. Gratwohl A, Hermans J, Goldman JM, et al. Risk assessment for patients with chronic myeloid leukaemia before allogeneic blood or marrow transplantation. Chronic Leukemia Working Party of the European Group



NCCN Guidelines Version 2.2022

Chronic Myeloid Leukemia

for Blood and Marrow Transplantation. *Lancet* 1998;352:1087-1092.

Available at: <http://www.ncbi.nlm.nih.gov/pubmed/9798583>.

274. Pavlu J, Kew AK, Taylor-Roberts B, et al. Optimizing patient selection for myeloablative allogeneic hematopoietic cell transplantation in chronic myeloid leukemia in chronic phase. *Blood* 2010;115:4018-4020. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/20304808>.

275. Horowitz MM, Rowlings PA, Passweg JR. Allogeneic bone marrow transplantation for CML: a report from the International Bone Marrow Transplant Registry. *Bone Marrow Transplant* 1996;17 Suppl 3:S5-6. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/8769690>.

276. Gratwohl A, Brand R, Apperley J, et al. Allogeneic hematopoietic stem cell transplantation for chronic myeloid leukemia in Europe 2006: transplant activity, long-term data and current results. An analysis by the Chronic Leukemia Working Party of the European Group for Blood and Marrow Transplantation (EBMT). *Haematologica* 2006;91:513-521. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/16533723>.

277. Goldman JM, Majhail NS, Klein JP, et al. Relapse and late mortality in 5-year survivors of myeloablative allogeneic hematopoietic cell transplantation for chronic myeloid leukemia in first chronic phase. *J Clin Oncol* 2010;28:1888-1895. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/20212247>.

278. Saussele S, Laussek M, Gratwohl A, et al. Allogeneic hematopoietic stem cell transplantation (allo SCT) for chronic myeloid leukemia in the imatinib era: evaluation of its impact within a subgroup of the randomized German CML Study IV. *Blood* 2010;115:1880-1885. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/19965667>.

279. Boehm A, Walcherberger B, Sperr WR, et al. Improved outcome in patients with chronic myeloid leukemia after allogeneic hematopoietic stem cell transplantation over the past 25 years: A single center experience. *Biol Blood Marrow Transplant* 2011 17:133-140. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/20601032>.

280. Khoury HJ, Kukreja M, Goldman JM, et al. Prognostic factors for outcomes in allogeneic transplantation for CML in the imatinib era: a CIBMTR analysis. *Bone Marrow Transplant* 2012;47:810-816. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/21986636>.

281. Delage R, Soiffer R, Dear K, Ritz J. Clinical significance of bcr-abl gene rearrangement detected by polymerase chain reaction after allogeneic bone marrow transplantation in chronic myelogenous leukemia. *Blood* 1991;78:2759-2767. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/1824268>.

282. Roth M, Antin J, Ash R, et al. Prognostic significance of Philadelphia chromosome-positive cells detected by the polymerase chain reaction after allogeneic bone marrow transplant for chronic myelogenous leukemia. *Blood* 1992;79:276-282. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/1728316>.

283. van Rhee F, Lin F, Cross NC, et al. Detection of residual leukaemia more than 10 years after allogeneic bone marrow transplantation for chronic myelogenous leukaemia. *Bone Marrow Transplant* 1994;14:609-612. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/7858536>.

284. Radich JP, Gehly G, Gooley T, et al. Polymerase chain reaction detection of the BCR-ABL fusion transcript after allogeneic marrow transplantation for chronic myeloid leukemia: results and implications in 346 patients. *Blood* 1995;85:2632-2638. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/7727789>.

285. Costello RT, Kirk J, Gabert J. Value of PCR analysis for long term survivors after allogeneic bone marrow transplant for chronic myelogenous leukemia: a comparative study. *Leuk Lymphoma* 1996;20:239-243. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/8624462>.

286. Mackinnon S, Barnett L, Heller G. Polymerase chain reaction is highly predictive of relapse in patients following T cell-depleted allogeneic bone marrow transplantation for chronic myeloid leukemia. *Bone Marrow Transplant* 1996;17:643-647. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/8722369>.

287. Olavarria E, Kanfer E, Szydlo R, et al. Early detection of BCR-ABL transcripts by quantitative reverse transcriptase-polymerase chain reaction predicts outcome after allogeneic stem cell transplantation for chronic myeloid leukemia. *Blood* 2001;97:1560-1565. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/11238091>.

288. Radich JP, Gooley T, Bryant E, et al. The significance of bcr-abl molecular detection in chronic myeloid leukemia patients "late," 18 months



NCCN Guidelines Version 2.2022

Chronic Myeloid Leukemia

or more after transplantation. Blood 2001;98:1701-1707. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/11535500>.

289. Kolb HJ, Schattenberg A, Goldman JM, et al. Graft-versus-leukemia effect of donor lymphocyte transfusions in marrow grafted patients. Blood 1995;86:2041-2050. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/7655033>.

290. Dazzi F, Szydlo RM, Cross NC, et al. Durability of responses following donor lymphocyte infusions for patients who relapse after allogeneic stem cell transplantation for chronic myeloid leukemia. Blood 2000;96:2712-2716. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/11023502>.

291. Luznik L, Fuchs EJ. Donor lymphocyte infusions to treat hematologic malignancies in relapse after allogeneic blood or marrow transplantation. Cancer Control 2002;9:123-137. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/11965233>.

292. Michallet AS, Nicolini F, Furst S, et al. Outcome and long-term follow-up of alloreactive donor lymphocyte infusions given for relapse after myeloablative allogeneic hematopoietic stem cell transplantations (HSCT). Bone Marrow Transplant 2005;35:601-608. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/15756285>.

293. Weissner M, Tischer J, Schnittger S, et al. A comparison of donor lymphocyte infusions or imatinib mesylate for patients with chronic myelogenous leukemia who have relapsed after allogeneic stem cell transplantation. Haematologica 2006;91:663-666. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/16627251>.

294. Chalandon Y, Passweg JR, Guglielmi C, et al. Early administration of donor lymphocyte infusions upon molecular relapse after allogeneic hematopoietic stem cell transplantation for chronic myeloid leukemia: a study by the Chronic Malignancies Working Party of the EBMT. Haematologica 2014;99:1492-1498. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/24997146>.

295. Schmidt S, Liu Y, Hu ZH, et al. The Role of Donor Lymphocyte Infusion (DLI) in Post-Hematopoietic Cell Transplant (HCT) Relapse for Chronic Myeloid Leukemia (CML) in the Tyrosine Kinase Inhibitor (TKI)

Era. Biol Blood Marrow Transplant 2020;26:1137-1143. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/32062061>.

296. Dazzi F, Szydlo RM, Craddock C, et al. Comparison of single-dose and escalating-dose regimens of donor lymphocyte infusion for relapse after allografting for chronic myeloid leukemia. Blood 2000;95:67-71. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/10607686>.

297. Shimoni A, Gajewski JA, Donato M, et al. Long-Term follow-up of recipients of CD8-depleted donor lymphocyte infusions for the treatment of chronic myelogenous leukemia relapsing after allogeneic progenitor cell transplantation. Biol Blood Marrow Transplant 2001;7:568-575. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/11760089>.

298. Gilleece MH, Dazzi F. Donor lymphocyte infusions for patients who relapse after allogeneic stem cell transplantation for chronic myeloid leukaemia. Leuk Lymphoma 2003;44:23-28. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/12691139>.

299. Posthuma EFM, Marijt EWF, Barge RMY, et al. Alpha-interferon with very-low-dose donor lymphocyte infusion for hematologic or cytogenetic relapse of chronic myeloid leukemia induces rapid and durable complete remissions and is associated with acceptable graft-versus-host disease. Biol Blood Marrow Transplant 2004;10:204-212. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/14993886>.

300. Simula MP, Marktel S, Fozza C, et al. Response to donor lymphocyte infusions for chronic myeloid leukemia is dose-dependent: the importance of escalating the cell dose to maximize therapeutic efficacy. Leukemia 2007;21:943-948. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/17361226>.

301. Kantarjian HM, O'Brien S, Cortes JE, et al. Imatinib mesylate therapy for relapse after allogeneic stem cell transplantation for chronic myelogenous leukemia. Blood 2002;100:1590-1595. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/12176876>.

302. Olavarria E, Ottmann OG, Deininger M, et al. Response to imatinib in patients who relapse after allogeneic stem cell transplantation for chronic myeloid leukemia. Leukemia 2003;17:1707-1712. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/12970768>.



NCCN Guidelines Version 2.2022

Chronic Myeloid Leukemia

303. Anderlini P, Sheth S, Hicks K, et al. Re: Imatinib mesylate administration in the first 100 days after stem cell transplantation. *Biol Blood Marrow Transplant* 2004;10:883-884. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/15570257>.

304. DeAngelo DJ, Hochberg EP, Alyea EP, et al. Extended follow-up of patients treated with imatinib mesylate (gleevec) for chronic myelogenous leukemia relapse after allogeneic transplantation: durable cytogenetic remission and conversion to complete donor chimerism without graft-versus-host disease. *Clin Cancer Res* 2004;10:5065-5071. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/15297408>.

305. Hess G, Bunjes D, Siegert W, et al. Sustained complete molecular remissions after treatment with imatinib-mesylate in patients with failure after allogeneic stem cell transplantation for chronic myelogenous leukemia: results of a prospective phase II open-label multicenter study. *J Clin Oncol* 2005;23:7583-7593. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/16234522>.

306. Palandri F, Amabile M, Rosti G, et al. Imatinib therapy for chronic myeloid leukemia patients who relapse after allogeneic stem cell transplantation: a molecular analysis. *Bone Marrow Transplant* 2007;39:189-191. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/17211436>.

307. Conchon M, Sanabani SS, Bendit I, et al. The use of imatinib mesylate as a lifesaving treatment of chronic myeloid leukemia relapse after bone marrow transplantation. *J Transplant* 2009;2009:357093-357093. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/20107580>.

308. Wright MP, Shepherd JD, Barnett MJ, et al. Response to tyrosine kinase inhibitor therapy in patients with chronic myelogenous leukemia relapsing in chronic and advanced phase following allogeneic hematopoietic stem cell transplantation. *Biol Blood Marrow Transplant* 2010;16:639-646. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/20005967>.

309. Breccia M, Cannella L, Stefanizzi C, et al. Efficacy of dasatinib in a chronic myeloid leukemia patient with disease molecular relapse and chronic GVHD after haploidentical BMT: an immunomodulatory effect?

Bone Marrow Transplant 2009;44:331-332. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/19219075>.

310. Klyuchnikov E, Schafhausen P, Kroger N, et al. Second-generation tyrosine kinase inhibitors in the post-transplant period in patients with chronic myeloid leukemia or Philadelphia-positive acute lymphoblastic leukemia. *Acta Haematol* 2009;122:6-10. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/19602874>.

311. Reinwald M, Schleyer E, Kiewe P, et al. Efficacy and pharmacologic data of second-generation tyrosine kinase inhibitor nilotinib in BCR-ABL-positive leukemia patients with central nervous system relapse after allogeneic stem cell transplantation. *Biomed Res Int* 2014;2014:637059. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/25025064>.

312. Shimoni A, Volchek Y, Koren-Michowitz M, et al. Phase 1/2 study of nilotinib prophylaxis after allogeneic stem cell transplantation in patients with advanced chronic myeloid leukemia or Philadelphia chromosome-positive acute lymphoblastic leukemia. *Cancer* 2015;121:863-871. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/25387866>.

313. Savani BN, Montero A, Kurlander R, et al. Imatinib synergizes with donor lymphocyte infusions to achieve rapid molecular remission of CML relapsing after allogeneic stem cell transplantation. *Bone Marrow Transplant* 2005;36:1009-1015. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/16205732>.

314. Shanavas M, Messner HA, Kamel-Reid S, et al. A comparison of long-term outcomes of donor lymphocyte infusions and tyrosine kinase inhibitors in patients with relapsed CML after allogeneic hematopoietic cell transplantation. *Clin Lymphoma Myeloma Leuk* 2014;14:87-92. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/24252361>.

315. Zeidner JF, Zahurak M, Rosner GL, et al. The evolution of treatment strategies for patients with chronic myeloid leukemia relapsing after allogeneic bone marrow transplant: can tyrosine kinase inhibitors replace donor lymphocyte infusions? *Leuk Lymphoma* 2015;56:128-134. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/24712979>.

316. Carpenter PA, Snyder DS, Flowers MED, et al. Prophylactic administration of imatinib after hematopoietic cell transplantation for high-



NCCN Guidelines Version 2.2022

Chronic Myeloid Leukemia

risk Philadelphia chromosome-positive leukemia. Blood 2007;109:2791-2793. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/17119111>.

317. Olavarria E, Siddique S, Griffiths MJ, et al. Posttransplantation imatinib as a strategy to postpone the requirement for immunotherapy in patients undergoing reduced-intensity allografts for chronic myeloid leukemia. Blood 2007;110:4614-4617. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/17881635>.

318. DeFilipp Z, Langston AA, Chen Z, et al. Does post-transplant maintenance therapy with tyrosine kinase inhibitors improve outcomes of patients with high-risk Philadelphia chromosome-positive leukemia? Clin Lymphoma Myeloma Leuk 2016;16:466-471 e461. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/27297665>.

319. Egan DN, Beppu L, Radich JP. Patients with Philadelphia-positive leukemia with BCR-ABL kinase mutations before allogeneic transplantation predominantly relapse with the same mutation. Biol Blood Marrow Transplant 2015;21:184-189. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/25300870>.

320. Oshima K, Kanda Y, Yamashita T, et al. Central nervous system relapse of leukemia after allogeneic hematopoietic stem cell transplantation. Biol Blood Marrow Transplant 2008;14:1100-1107. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/18804039>.

321. Fuchs M, Reinhofer M, Ragooschke-Schumm A, et al. Isolated central nervous system relapse of chronic myeloid leukemia after allogeneic hematopoietic stem cell transplantation. BMC Blood Disord 2012;12:9. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/22871019>.

322. Ocheni S, Iwanski GB, Schafhausen P, et al. Characterisation of extramedullary relapse in patients with chronic myeloid leukemia in advanced disease after allogeneic stem cell transplantation. Leuk Lymphoma 2009;50:551-558. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/19373652>.

323. Nishimoto M, Nakamae H, Koh KR, et al. Dasatinib maintenance therapy after allogeneic hematopoietic stem cell transplantation for an isolated central nervous system blast crisis in chronic myelogenous leukemia. Acta Haematol 2013;130:111-114. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/23548721>.

324. Massimino M, Stella S, Tirro E, et al. Non ABL-directed inhibitors as alternative treatment strategies for chronic myeloid leukemia. Mol Cancer 2018;17:56. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/29455672>.

325. Annunziata M, Bonifacio M, Breccia M, et al. Current Strategies and Future Directions to Achieve Deep Molecular Response and Treatment-Free Remission in Chronic Myeloid Leukemia. Front Oncol 2020;10:883. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/32582549>.

326. Clark RE, Polydoros F, Apperley JF, et al. De-escalation of tyrosine kinase inhibitor therapy before complete treatment discontinuation in patients with chronic myeloid leukaemia (DESTINY): a non-randomised, phase 2 trial. Lancet Haematol 2019;6:e375-e383. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/31201085>.

327. Westerweel PE, Te Boekhorst PAW, Levin MD, Cornelissen JJ. New Approaches and Treatment Combinations for the Management of Chronic Myeloid Leukemia. Front Oncol 2019;9:665. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/31448223>.

328. Hoffmann VS, Baccarani M, Hasford J, et al. The EUTOS population-based registry: incidence and clinical characteristics of 2904 CML patients in 20 European Countries. Leukemia 2015;29:1336-1343. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/25783795>.

329. Ramasamy K, Hayden J, Lim Z, et al. Successful pregnancies involving men with chronic myeloid leukaemia on imatinib therapy. Br J Haematol 2007;137:374-375. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/17408403>.

330. Breccia M, Cannella L, Montefusco E, et al. Male patients with chronic myeloid leukemia treated with imatinib involved in healthy pregnancies: report of five cases. Leuk Res 2008;32:519-520. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/17804066>.

331. Oweini H, Otrrock ZK, Mahfouz RAR, Bazarbachi A. Successful pregnancy involving a man with chronic myeloid leukemia on dasatinib. Arch Gynecol Obstet 2010. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/20473616>.

332. Ghalaut VS, Prakash G, Bansal P, et al. Effect of imatinib on male reproductive hormones in BCR-ABL positive CML patients: A preliminary



NCCN Guidelines Version 2.2022

Chronic Myeloid Leukemia

report. J Oncol Pharm Pract 2014;20:243-248. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/23966360>.

333. Alizadeh H, Jaafar H, Rajnics P, et al. Outcome of pregnancy in chronic myeloid leukaemia patients treated with tyrosine kinase inhibitors: short report from a single centre. Leuk Res 2015;39:47-51. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/25455655>.

334. Pye SM, Cortes J, Ault P, et al. The effects of imatinib on pregnancy outcome. Blood 2008;111:5505-5508. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/18322153>.

335. Cortes JE, Abruzzese E, Chelysheva E, et al. The impact of dasatinib on pregnancy outcomes. Am J Hematol 2015;90:1111-1115. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/26348106>.

336. Barkoulas T, Hall PD. Experience with dasatinib and nilotinib use in pregnancy. J Oncol Pharm Pract 2018;24:121-128. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/29284357>.

337. Salem W, Li K, Krapp C, et al. Imatinib treatments have long-term impact on placentation and embryo survival. Sci Rep 2019;9:2535. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/30796277>.

338. Madabhavi I, Sarkar M, Modi M, Kadakol N. Pregnancy Outcomes in Chronic Myeloid Leukemia: A Single Center Experience. J Glob Oncol 2019;5:1-11. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/31584851>.

339. Ault P, Kantarjian H, O'Brien S, et al. Pregnancy among patients with chronic myeloid leukemia treated with imatinib. J Clin Oncol 2006;24:1204-1208. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/16446320>.

340. Kuwabara A, Babb A, Ibrahim A, et al. Poor outcome after reintroduction of imatinib in patients with chronic myeloid leukemia who interrupt therapy on account of pregnancy without having achieved an optimal response. Blood 2010;116:1014-1016. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/20705771>.

341. Lasica M, Willcox A, Burbury K, et al. The effect of tyrosine kinase inhibitor interruption and interferon use on pregnancy outcomes and long-

term disease control in chronic myeloid leukemia. Leuk Lymphoma 2019;1-7. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/30632843>.

342. Stella S, Tirro E, Massimino M, et al. Successful Management of a Pregnant Patient With Chronic Myeloid Leukemia Receiving Standard Dose Imatinib. In Vivo 2019;33:1593-1598. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/31471409>.

343. Haggstrom J, Adriansson M, Hybbinette T, et al. Two cases of CML treated with alpha-interferon during second and third trimester of pregnancy with analysis of the drug in the new-born immediately postpartum. Eur J Haematol 1996;57:101-102. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/8698119>.

344. Kuroiwa M, Gondo H, Ashida K, et al. Interferon-alpha therapy for chronic myelogenous leukemia during pregnancy. Am J Hematol 1998;59:101-102. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/9723590>.

345. Lipton JH, Derzko CM, Curtis J. Alpha-interferon and pregnancy in a patient with CML. Hematol Oncol 1996;14:119-122. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/9119356>.

346. Baykal C, Zengin N, Coskun F, et al. Use of hydroxyurea and alpha-interferon in chronic myeloid leukemia during pregnancy: a case report. Eur J Gynaecol Oncol 2000;21:89-90. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/10726630>.

347. Thauvin-Robinet C, Maingueneau C, Robert E, et al. Exposure to hydroxyurea during pregnancy: a case series. Leukemia 2001;15:1309-1311. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/11480579>.

348. Fadilah SA, Ahmad-Zailani H, Soon-Keng C, Norlaila M. Successful treatment of chronic myeloid leukemia during pregnancy with hydroxyurea. Leukemia 2002;16:1202-1203. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/12040456>.

349. Al Bahar S, Pandita R, Nath SV. Pregnancy in chronic myeloid leukemia patients treated with alpha interferon. Int J Gynaecol Obstet 2004;85:281-282. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/15145269>.



NCCN Guidelines Version 2.2022

Chronic Myeloid Leukemia

350. Koh LP, Kanagalingam D. Pregnancies in patients with chronic myeloid leukemia in the era of imatinib. *Int J Hematol* 2006;84:459-462. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/17189230>.

351. Balsat M, Etienne M, Elhamri M, et al. Successful pregnancies in patients with BCR-ABL-positive leukemias treated with interferon-alpha therapy during the tyrosine kinase inhibitors era. *Eur J Haematol* 2018;101:774-780. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/30179268>.

352. Burchert A, Muller MC, Kostrewa P, et al. Sustained molecular response with interferon alfa maintenance after induction therapy with imatinib plus interferon alfa in patients with chronic myeloid leukemia. *J Clin Oncol* 2010;28:1429-1435. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/20142590>.

353. Abruzzese E, Turkina AG, Apperley JF, et al. Pregnancy Management in CML Patients: To Treat or Not to Treat? Report of 224 Outcomes of the European Leukemia Net (ELN) Database. *Blood* 2019;134:498-498. Available at: <https://doi.org/10.1182/blood-2019-124430>.

354. Beauverd Y, Radia D, Cargo C, et al. Pegylated interferon alpha-2a for essential thrombocythemia during pregnancy: outcome and safety. A case series. *Haematologica* 2016;101:e182-184. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/26819057>.

355. Ali R, Ozkalemkas F, Ozkocaman V, et al. Successful pregnancy and delivery in a patient with chronic myelogenous leukemia (CML), and management of CML with leukapheresis during pregnancy: a case report and review of the literature. *Jpn J Clin Oncol* 2004;34:215-217. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/15121759>.

356. Palani R, Milojkovic D, Apperley JF. Managing pregnancy in chronic myeloid leukaemia. *Ann Hematol* 2015;94 Suppl 2:S167-176. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/25814083>.

357. Staley EM, Simmons SC, Feldman AZ, et al. Management of chronic myeloid leukemia in the setting of pregnancy: when is leukocytapheresis appropriate? A case report and review of the literature. *Transfusion* 2018;58:456-460. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/29230832>.

358. James AH, Brancazio LR, Price T. Aspirin and reproductive outcomes. *Obstet Gynecol Surv* 2008;63:49-57. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/18081940>.

359. Deruelle P, Coulon C. The use of low-molecular-weight heparins in pregnancy--how safe are they? *Curr Opin Obstet Gynecol* 2007;19:573-577. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/18007136>.

360. Russell MA, Carpenter MW, Akhtar MS, et al. Imatinib mesylate and metabolite concentrations in maternal blood, umbilical cord blood, placenta and breast milk. *J Perinatol* 2007;27:241-243. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/17377606>.

361. Ali R, Ozkalemkas F, Kimya Y, et al. Imatinib use during pregnancy and breast feeding: a case report and review of the literature. *Arch Gynecol Obstet* 2009;280:169-175. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/19083009>.

362. Drugs and Lactation Database (LactMed) [Internet]. Bethesda (MD): National Library of Medicine (US); 2006-. Available at: <https://www.ncbi.nlm.nih.gov/books/NBK501922/>.

363. Chelysheva E, Aleshin S, Polushkina E, et al. Breastfeeding in Patients with Chronic Myeloid Leukaemia: Case Series with Measurements of Drug Concentrations in Maternal Milk and Literature Review. *Mediterr J Hematol Infect Dis* 2018;10:e2018027. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/29755704>.

364. Abruzzese E, Trawinska MM, Perrotti AP, De Fabritiis P. Tyrosine kinase inhibitors and pregnancy. *Mediterr J Hematol Infect Dis* 2014;6:e2014028. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/24804001>.

365. Hijiya N, Suttorp M. How I treat chronic myeloid leukemia in children and adolescents. *Blood* 2019;133:2374-2384. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/30917954>.

366. Suttorp M, Schulze P, Glauche I, et al. Front-line imatinib treatment in children and adolescents with chronic myeloid leukemia: results from a phase III trial. *Leukemia* 2018;32:1657-1669. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/29925908>.



NCCN Guidelines Version 2.2022

Chronic Myeloid Leukemia

367. Gore L, Kearns PR, de Martino ML, et al. Dasatinib in Pediatric Patients With Chronic Myeloid Leukemia in Chronic Phase: Results From a Phase II Trial. *J Clin Oncol* 2018;36:1330-1338. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/29498925>.

368. Hijjiya N, Maschan A, Rizzari C, et al. Phase 2 study of nilotinib in pediatric patients with Philadelphia chromosome-positive chronic myeloid leukemia. *Blood* 2019;134:2036-2045. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/31511239>.

369. Millot F, Guilhot J, Nelken B, et al. Imatinib mesylate is effective in children with chronic myelogenous leukemia in late chronic and advanced phase and in relapse after stem cell transplantation. *Leukemia* 2006;20:187-192. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/16341042>.

370. Champagne MA, Fu CH, Chang M, et al. Higher dose imatinib for children with de novo chronic phase chronic myelogenous leukemia: a report from the Children's Oncology Group. *Pediatr Blood Cancer* 2011;57:56-62. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/21465636>.

371. Giona F, Putti MC, Micalizzi C, et al. Long-term results of high-dose imatinib in children and adolescents with chronic myeloid leukaemia in chronic phase: the Italian experience. *Br J Haematol* 2015;170:398-407. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/25891192>.

372. Nickel RS, Daves M, Keller F. Treatment of an adolescent with chronic myeloid leukemia and the T315I mutation with ponatinib. *Pediatr Blood Cancer* 2015;62:2050-2051. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/25939962>.

373. Gurrea Salas D, Glauche I, Tauer JT, et al. Can prognostic scoring systems for chronic myeloid leukemia as established in adults be applied to pediatric patients? *Ann Hematol* 2015;94:1363-1371. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/25894600>.

374. Millot F, Guilhot J, Suttorp M, et al. Prognostic discrimination based on the EUTOS long-term survival score within the International Registry for Chronic Myeloid Leukemia in children and adolescents. *Haematologica* 2017;102:1704-1708. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/28838993>.

375. Samis J, Lee P, Zimmerman D, et al. Recognizing endocrinopathies associated with tyrosine kinase inhibitor therapy in children with chronic myelogenous leukemia. *Pediatr Blood Cancer* 2016;63:1332-1338. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/27100618>.

376. Suttorp M, Millot F. Treatment of pediatric chronic myeloid leukemia in the year 2010: use of tyrosine kinase inhibitors and stem-cell transplantation. *Hematology Am Soc Hematol Educ Program* 2010;2010:368-376. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/21239821>.

377. Shima H, Tokuyama M, Tanizawa A, et al. Distinct impact of imatinib on growth at prepubertal and pubertal ages of children with chronic myeloid leukemia. *J Pediatr* 2011;159:676-681. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/21592517>.

378. Bansal D, Shava U, Varma N, et al. Imatinib has adverse effect on growth in children with chronic myeloid leukemia. *Pediatr Blood Cancer* 2012;59:481-484. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/22052850>.

379. Rastogi MV, Stork L, Druker B, et al. Imatinib mesylate causes growth deceleration in pediatric patients with chronic myelogenous leukemia. *Pediatr Blood Cancer* 2012;59:840-845. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/22378641>.

380. Narayanan KR, Bansal D, Walia R, et al. Growth failure in children with chronic myeloid leukemia receiving imatinib is due to disruption of GH/IGF-1 axis. *Pediatr Blood Cancer* 2013;60:1148-1153. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/23322583>.

381. Boddu D, Thankamony P, Guruprasad CS, et al. Effect of imatinib on growth in children with chronic myeloid leukemia. *Pediatr Hematol Oncol* 2019;36:189-197. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/31298597>.

382. Sabnis HS, Keenum C, Lewis RW, et al. Growth disturbances in children and adolescents receiving long-term tyrosine kinase inhibitor therapy for Chronic Myeloid Leukaemia or Philadelphia Chromosome-positive Acute Lymphoblastic Leukaemia. *Br J Haematol* 2019;185:795-799. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/30407613>.



NCCN Guidelines Version 2.2022

Chronic Myeloid Leukemia

383. de Bruijn CMA, Millot F, Suttrop M, et al. Discontinuation of imatinib in children with chronic myeloid leukaemia in sustained deep molecular remission: results of the STOP IMAPED study. *Br J Haematol* 2019. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/30843196>.

384. Millot F, Claviez A, Leverger G, et al. Imatinib cessation in children and adolescents with chronic myeloid leukemia in chronic phase. *Pediatr Blood Cancer* 2014;61:355-357. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/24106110>.

385. de Lavallade H, Khoder A, Hart M, et al. Tyrosine kinase inhibitors impair B-cell immune responses in CML through off-target inhibition of kinases important for cell signaling. *Blood* 2013;122:227-238. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/23719297>.

386. de Lavallade H, Garland P, Sekine T, et al. Repeated vaccination is required to optimize seroprotection against H1N1 in the immunocompromised host. *Haematologica* 2011;96:307-314. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/20971824>.

387. Luthy KE, Tiedeman ME, Beckstrand RL, Mills DA. Safety of live-virus vaccines for children with immune deficiency. *J Am Acad Nurse Pract* 2006;18:494-503. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/16999715>.

388. Hughes TP, Mauro MJ, Cortes JE, et al. Asciminib in Chronic Myeloid Leukemia after ABL Kinase Inhibitor Failure. *N Engl J Med* 2019;381:2315-2326. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/31826340>.

389. Do YR, Kwak JY, Kim JA, et al. Long-term data from a phase 3 study of radotinib versus imatinib in patients with newly diagnosed, chronic myeloid leukaemia in the chronic phase (RERISE). *Br J Haematol* 2020;189:303-312. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/32012231>.

390. Kim SH, Menon H, Jootar S, et al. Efficacy and safety of radotinib in chronic phase chronic myeloid leukemia patients with resistance or intolerance to BCR-ABL1 tyrosine kinase inhibitors. *Haematologica* 2014;99:1191-1196. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/24705186>.

391. Cortes J, Jabbour E, Daley GQ, et al. Phase 1 study of lonafermin (SCH 66336) and imatinib mesylate in patients with chronic myeloid leukemia who have failed prior single-agent therapy with imatinib. *Cancer* 2007;110:1295-1302. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/17623836>.

392. Cortes J, Quintas-Cardama A, Garcia-Manero G, et al. Phase 1 study of tipifarnib in combination with imatinib for patients with chronic myelogenous leukemia in chronic phase after imatinib failure. *Cancer* 2007;110:2000-2006. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/17849425>.

393. Borthakur G, Dombret H, Schafhausen P, et al. A phase I study of danusertib (PHA-739358) in adult patients with accelerated or blastic phase chronic myeloid leukemia and Philadelphia chromosome-positive acute lymphoblastic leukemia resistant or intolerant to imatinib and/or other second generation c-ABL therapy. *Haematologica* 2015;100:898-904. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/25887498>.

394. Seymour JF, Kim DW, Rubin E, et al. A phase 2 study of MK-0457 in patients with BCR-ABL T315I mutant chronic myelogenous leukemia and Philadelphia chromosome-positive acute lymphoblastic leukemia. *Blood Cancer J* 2014;4:e238. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/25127392>.

395. Sweet K, Hazlehurst L, Sahakian E, et al. A phase I clinical trial of ruxolitinib in combination with nilotinib in chronic myeloid leukemia patients with molecular evidence of disease. *Leuk Res* 2018;74:89-96. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/30340199>.